

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 May 2003 (08.05.2003)

PCT

(10) International Publication Number  
**WO 03/037860 A2**

(51) International Patent Classification<sup>7</sup>: **C07D**

**FAN, Junhua** [US/US]; 7528 Charmant Drive, San Diego, CA 92122 (US).

(21) International Application Number: PCT/US02/35069

(22) International Filing Date: 30 October 2002 (30.10.2002)

(74) Agent: **KREUSSER, Edward O.**; BROBECK PHLEGER & HARRISON LLP, 12390 El Camino Real, San Diego, CA 92130-2081 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/335,391 30 October 2001 (30.10.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*):  
**CONFORMA THERAPEUTICS CORPORATION** [US/US]; 9393 Towne Centre Drive, Suite 240, San Diego, CA 92121 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **KASIBHATLA, Srinivas Rao** [US/US]; 13073 Dressage Lane, San Diego, CA 92130 (US). **HONG, Kevin** [US/US]; 5734 Redwood Street, San Diego, CA 92105 (US). **ZHANG, Lin** [US/US]; 12455 Brickellia Street, San Diego, CA 92129 (US). **BIAMONTE, Marco Antonio** [US/US]; 5350 Toscana Way, Apt. 216, San Diego, CA 92122 (US). **BOEHM, Marcus F.** [US/US]; 38833 Lamont Street, #4E, San Diego, CA 92109 (US). **SHI, Jiandong** [US/US]; 8540 Costa Verde Blvd., San Diego, CA 92122 (US).

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PURINE ANALOGS HAVING HSP90-INHIBITING ACTIVITY

(57) Abstract: Novel purine compounds and tautomers and pharmaceutically acceptable salts thereof are described, as are pharmaceutical compositions comprising the same, complexes comprising the same, e.g., HSP90 complexes, and methods of using the same.



WO 03/037860 A2

## PURINE ANALOGS HAVING HSP90-INHIBITING ACTIVITY

## RELATED APPLICATIONS

This application claims priority to and incorporates by reference in its entirety Kasibhatla et al., United States Provisional Patent Application Serial Number 60/335,391,  
5 entitled PURINE ANALOGS HAVING HSP90-INHIBITING ACTIVITY, filed October 30, 2001.

## FIELD OF THE INVENTION

The invention relates in general to purine analogs and their use in inhibiting heat shock protein 90's (HSP90's) to thereby treat or prevent HSP90-dependent diseases, e.g.,  
10 proliferative disorders such as breast cancer.

## BACKGROUND

The following description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed inventions, or that any  
15 publication specifically or implicitly referenced is prior art.

HSP90s are ubiquitous chaperone proteins that are involved in folding, activation and assembly of a wide range of proteins, including key proteins involved in signal transduction, cell cycle control and transcriptional regulation. Researchers have reported that HSP90 chaperone proteins are associated with important signaling proteins, such as  
20 steroid hormone receptors and protein kinases, including, e.g., Raf-1, EGFR, v-Src family kinases, Cdk4, and ErbB-2 ( Buchner J., 1999, TIBS, 24:136-141; Stepanova, L. et al., 1996, Genes Dev. 10:1491-502; Dai, K. et al., 1996, J. Biol. Chem. 271:22030-4). Studies further indicate that certain co-chaperones, e.g., Hsp70, p60/Hop/Sti1, Hip, Bag1, HSP40/Hdj2/Hsj1, immunophilins, p23, and p50, may assist HSP90 in its function (see,  
25 e.g., Caplan, A., Trends in Cell Biol., 9: 262-68 (1999).

Ansamycin antibiotics, e.g., herbimycin A (HA), geldanamycin (GM), and 17-AAG are thought to exert their anticancerous effects by tight binding of the N-terminus pocket of HSP90, thereby destabilizing substrates that normally interact with HSP90

(Stebbins, C. et al., 1997, *Cell*, 89:239-250). This pocket is highly conserved and has weak homology to the ATP-binding site of DNA gyrase (Stebbins, C. et al., supra; Grenert, J.P. et al., 1997, *J. Biol. Chem.*, 272:23843-50). Further, ATP and ADP have both been shown to bind this pocket with low affinity and to have weak ATPase activity (Proromou, C. et al., 1997, *Cell*, 90: 65-75; Panaretou, B. et al., 1998, *EMBO J.*, 17: 4829-36). In vitro and in vivo studies have demonstrated that occupancy of this N-terminal pocket by ansamycins and other HSP90 inhibitors alters HSP90 function and inhibits protein folding. At high concentrations, ansamycins and other HSP90 inhibitors have been shown to prevent binding of protein substrates to HSP90 (Scheibel, T., H. et al., 1999, *Proc. Natl. Acad. Sci. U S A* 96:1297-302; Schulte, T. W. et al., 1995, *J. Biol. Chem.* 270:24585-8; Whitesell, L., et al., 1994, *Proc. Natl. Acad. Sci. U S A* 91:8324-8328). Ansamycins have also been demonstrated to inhibit the ATP-dependent release of chaperone-associated protein substrates (Schneider, C., L. et al., 1996, *Proc. Natl. Acad. Sci. U S A*, 93:14536-41; Sepp-Lorenzino et al., 1995, *J. Biol. Chem.* 270:16580-16587). In either event, the substrates are degraded by a ubiquitin-dependent process in the proteasome (Schneider, C., L., supra; Sepp-Lorenzino, L., et al., 1995, *J. Biol. Chem.*, 270:16580-16587; Whitesell, L. et al., 1994, *Proc. Natl. Acad. Sci. USA*, 91: 8324-8328).

HSP90 substrate destabilization occurs in tumor and non-transformed cells alike and has been shown to be especially effective on a subset of signaling regulators, e.g., Raf (Schulte, T. W. et al., 1997, *Biochem. Biophys. Res. Commun.* 239:655-9; Schulte, T. W., et al., 1995, *J. Biol. Chem.* 270:24585-8), nuclear steroid receptors (Segnitz, B., and U. Gehring. 1997, *J. Biol. Chem.* 272:18694-18701; Smith, D. F. et al., 1995, *Mol. Cell. Biol.* 15:6804-12 ), v-src (Whitesell, L., et al., 1994, *Proc. Natl. Acad. Sci. U S A* 91:8324-8328) and certain transmembrane tyrosine kinases (Sepp-Lorenzino, L. et al., 1995, *J. Biol. Chem.* 270:16580-16587) such as EGF receptor (EGFR) and Her2/Neu (Hartmann, F., et al., 1997, *Int. J. Cancer* 70:221-9; Miller, P. et al., 1994, *Cancer Res.* 54:2724-2730; Mimnaugh, E. G., et al., 1996, *J. Biol. Chem.* 271:22796-801; Schnur, R. et al., 1995, *J. Med. Chem.* 38:3806-3812), CDK4, and mutant p53. Erlichman et al., *Proc. AACR* (2001), 42, abstract 4474. The ansamycin-induced loss of these proteins leads to the selective disruption of certain regulatory pathways and results in growth arrest at specific phases of the cell cycle (Muisse-Heimericks, R. C. et al., 1998, *J. Biol. Chem.*

273:29864-72), and apoptosis, and/or differentiation of cells so treated (Vasilevskaya, A. et al., 1999, Cancer Res., 59:3935-40).

In addition to anti-cancer and antitumorigenic activity, HSP90 inhibitors have also been implicated in a wide variety of other utilities, including use as anti-inflammation agents, anti-infectious disease agents, agents for treating autoimmunity, agents for treating ischemia, and agents useful in promoting nerve regeneration (*See, e.g.*, Rosen et al., WO 02/09696; PCT/US01/23640; Degranco et al., WO 99/51223; PCT/US99/07242; Gold, U.S. Patent 6,210,974 B1). There are reports in the literature that fibrogenetic disorders including but not limited to scleroderma, polymyositis, systemic lupus, rheumatoid arthritis, liver cirrhosis, keloid formation, interstitial nephritis, and pulmonary fibrosis may be treatable. (Strehlow, WO 02/02123; PCT/US01/20578).


Ansamycins and other HSP90 inhibitors thus hold great promise for the treatment and/or prevention of many types of disorders. However, their relative insolubility makes them difficult to formulate and administer, and they are not easily synthesized and currently must, at least in part, be generated through fermentation. Further, the dose limiting toxicity of ansamycins is hepatic. Despite the potential of ansamycins, alternative HSP90 inhibitors are therefore needed.

Recently, Chiosis et al. described the design and synthesis of purine analogs that mimic geldanamycin and other ansamycins in their ability to bind the ATP binding pocket of, and thus inhibit, HSP90. *See* International Patent Application PCT/US01/46303 (WO 02/36075; Chemistry & Biology 8:289-299 (2001)). The specific compounds that Chiosis et al. described included a trimethoxybenzyl entity substituted at positions 3,4, and 5. Using gel-binding assays, these were shown to bind HSP90 approximately 20-fold less avidly than the benzoquinone ansamycin, 17-AAG. Chiosis et al. did not attempt a quinone mimic for the methoxybenzyl entity, speculating that to do so would lead to hepatotoxicity. *Id.*, pg. 290, col. 1, ¶ 4. Nor did Chiosis et al. teach, suggest, or otherwise report the use of sulfides, sulfoxides, and sulfones as described herein.


## SUMMARY OF THE INVENTION

Applicants herein describe a set of purine-based compounds that have utility in inhibiting HSP90 and diseases that are HSP90-dependent, e.g., a variety of carcinomas, such as melanoma, breast cancer, etc.


5 In one embodiment, the purine or purine analog has structure I, II, III, or IV:




I



II



III



IV

wherein A is selected from  $\text{NR}^1_2$ ,  $\text{NHSO}_2\text{R}^2$ ,  $\text{NR}^1\text{NR}^1_2$ ,  $\text{NR}^1\text{OR}^4$ ,  $\text{OR}^3$ ,  $\text{SR}^3$ , optionally substituted lower alkyl,  $\text{C(O)N(R}^4)_2$ , guanidine, amidine, H, halogen, CN,  $\text{N}_3$  and perhaloalkyl;


10 wherein X is a 1 carbon, 2 carbon, or 3 carbon optionally substituted structure (C1-C3), or else  $\text{NR}^1$ , S,  $\text{S(O)}$ ,  $\text{S(O)}_2$ , O, or  $\text{C(O)}$ . For carbon linkers having more than 1 carbon, these may have single, double, or triple bonds between them.

15 wherein Y is selected from the group consisting of H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted alicyclic, optionally substituted araalkyl, optionally substituted aryloxyalkyl, optionally substituted alkoxyalkyl, alkylaminoalkyl,

alkylcarbonylaminoalkyl  $(-(CH_2)_n-C(O)-NR-(CH_2)_n)$ , alkylcarbonyloxyalkyl  $(-(CH_2)_n-C(O)-O-(CH_2)_n)$ , optionally substituted heterocyclic, hydroxyalkyl, haloalkyl, perhaloalkyl,  $C(O)R^2$ ,  $S(O)_2R^2$ ,  $C(O)NHR^2$ , and  $C(O)OR^2$ ;

5 wherein Z is selected from the group consisting of H, halogen, CN,  $OR^3$ ,  $SR^3$ , perhaloalkyl, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted alicyclic, optionally substituted araalkyl, optionally substituted aryloxyalkyl, optionally substituted alkoxyalkyl, optionally substituted heterocyclic,  $C(O)R^2$ ,  $-S(O)_2R^2$ ,  $NHOR^4$ , and  $C(O)NR^4_2$ ;

10 Q is selected from the group consisting of alkyl, cycloalkyl, arylalkyl, aryl, heteroaryl, and heterocyclic, all optionally substituted; e. g.,



$L = O, S, NR^1$

15  $R^1$  is independently selected from H, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heteroalkyl, optionally substituted aryl, optionally substituted heterocyclic,  $C(O)R^2$ ,  $-C(O)OR^2$ ,  $C(O)NR^4_2$ ,  $C(S)OR^2$ ,  $C(S)NR^4_2$ ,  $PO_3R_4$ , and  $SO_2R^2$ ;

$R^2$  is selected from alkyl, heteroalkyl, cycloalkyl, heterocyclic, heteroaryl, and aryl, all optionally substituted;

$R^3$  is selected from H, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heteroalkyl, optionally substituted aryl, optionally substituted heterocyclic and  $C(O)NR^4_2$ ; and


wherein  $R^4$  is selected from either H or from alkyl, cycloalkyl, heteroalkyl, aryl, and heterocyclic, all optionally substituted;

wherein  $R^5$  is selected from H, OH, and optionally substituted alkyl; and

wherein  $R^6$  is independently selected from H, optionally substituted alkyl,  $OR^3$ ,  $SR^3$ ,  $NHR^3$ ,  $C(O)R^5$ ,  $NO_2$ , CN, halogen, and  $S(O)_2R^2$ . In some embodiments, the alkyl, alkenyl, and alkynyl substituent are 1 to 8 carbon atoms in length, more preferably 1 to 6 carbon atoms in length, and optionally substituted.

The subscript "n" can be 1 to 10 inclusive, with 3 or less being preferred.

Other compounds of the invention are based on the following formula, having illustrative species as described in Table 1:



**Table 1**

No.	Ex.	W	X	Y	Z
1	3.2	2,5-dimethoxy	$CH_2$	pent-4-ynyl - $(CH_2)_3CCH$	H
2		2,3,5-trimethoxy	$CH_2$	pent-4-ynyl	H
3		3,4,5-trimethoxy	$CH_2$	pent-4-ynyl	H
4	4.7	2-iodo,5-methoxy	$CH_2$	pent-4-ynyl	H
5		2-bromo,5-methoxy	$CH_2$	pent-4-ynyl	H

No.	Ex.	W	X	Y	Z
6		2-chloro,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
7		2,4-diiodo,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
8		2,5-diiodo	CH <sub>2</sub>	pent-4-ynyl	H
9		2,4-diiodo	CH <sub>2</sub>	pent-4-ynyl	H
10		2-iodo,5-SCH <sub>3</sub>	CH <sub>2</sub>	pent-4-ynyl	H
11		2-iodo,5-ethyl	CH <sub>2</sub>	pent-4-ynyl	H
12		2-iodo,5-propyl	CH <sub>2</sub>	pent-4-ynyl	H
13		2-chloro,5-SCH <sub>3</sub>	CH <sub>2</sub>	pent-4-ynyl	H
14		2-chloro,5-ethyl	CH <sub>2</sub>	pent-4-ynyl	H
15		2-chloro,5-propyl	CH <sub>2</sub>	pent-4-ynyl	H
16		2,5-SCH <sub>3</sub>	CH <sub>2</sub>	pent-4-ynyl	H
17		2-iodo, 4-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
18		2-iodo, 3-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
19		2-iodo, 6-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
20		2-Chloro, 3-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
21		2-Chloro, 4-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
22		2-chloro-3,4,5-trimethoxy	CH <sub>2</sub>	pent-4-ynyl	H
23		2,3-diiodo,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
24		2,5-dichloro	CH <sub>2</sub>	pent-4-ynyl	H
25		2,5-dibromo	CH <sub>2</sub>	pent-4-ynyl	H
26		2-Iodo,4-chloro,5-	CH <sub>2</sub>	pent-4-ynyl	H



No.	Ex.	W	X	Y	Z
		methoxy			
27		2-iodo,4-bromo,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
28		2,5-dimethoxy	CH <sub>2</sub>	pent-4-ynyl	F
29		2,3,5-trimethoxy	CH <sub>2</sub>	pent-4-ynyl	F
30		3,4,5-trimethoxy	CH <sub>2</sub>	pent-4-ynyl	F
31	4.10	2-iodo,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
32		2-bromo,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
33		2-chloro,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
34		2,4-diiodo,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
35		2,5-diiodo	CH <sub>2</sub>	pent-4-ynyl	F
36		2,4-diiodo	CH <sub>2</sub>	pent-4-ynyl	F
37		2-iodo,5-SCH <sub>3</sub>	CH <sub>2</sub>	pent-4-ynyl	F
38		2-iodo,5-ethyl	CH <sub>2</sub>	pent-4-ynyl	F
39		2-iodo,5-propyl	CH <sub>2</sub>	pent-4-ynyl	F
40		2-chloro,5-SCH <sub>3</sub>	CH <sub>2</sub>	pent-4-ynyl	F
41		2-chloro,5-ethyl	CH <sub>2</sub>	pent-4-ynyl	F
42		2-chloro,5-propyl	CH <sub>2</sub>	pent-4-ynyl	F
43		2,5-SCH <sub>3</sub>	CH <sub>2</sub>	pent-4-ynyl	F
44		2-iodo, 4-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
45		2-iodo, 3-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
46		2-iodo, 6-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
47		2-Chloro, 3-fluoro, 5-	CH <sub>2</sub>	pent-4-ynyl	F

No.	Ex.	W	X	Y	Z
		methoxy			
48		2-Chloro, 4-fluoro, 5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
49		2,3,4,5-tetraiodo	CH <sub>2</sub>	pent-4-ynyl	F
50		2,3-diiodo,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
51		2,5-dichloro	CH <sub>2</sub>	pent-4-ynyl	F
52		2,5-dibromo	CH <sub>2</sub>	pent-4-ynyl	F
53		2-Iodo,4-chloro,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
54		2-iodo,4-bromo,5-methoxy	CH <sub>2</sub>	pent-4-ynyl	F
55	3.4	2,5-dimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl -(CH <sub>2</sub> ) <sub>2</sub> CHCCMe <sub>2</sub>	H
56		2,3,5-trimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
57		3,4,5-trimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
58	4.8	2-iodo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
59		2-bromo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
60		2-chloro,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
61		2,4-diiodo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
62		2,5-diiodo	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
63		2,4-diiodo	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
64		2-iodo,5-SCH <sub>3</sub>	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
65		2-iodo,5-ethyl	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
66		2-iodo,5-propyl	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
67		2-chloro,5-SCH <sub>3</sub>	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
68		2-chloro,5-ethyl	CH <sub>2</sub>	4-methyl-pent-3-enyl	H

No.	Ex.	W	X	Y	Z
69		2-chloro,5-propyl	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
70		2,5-SCH <sub>3</sub>	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
71		2-iodo, 4-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
72		2-iodo, 3-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
73		2-iodo, 6-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
74		2-Chloro, 3-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
75		2-Chloro, 4-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
76		2-chloro-3,4,5-trimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
77		2,3-diiodo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
78		2,5-dichloro	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
79		2,5-dibromo	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
80		2-iodo,4-chloro,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
81		2-iodo,4-bromo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
82		2,5-dimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
83		2,3,5-trimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
84		3,4,5-trimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
85	4.9	2-iodo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
86		2-bromo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
87		2-chloro,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
88		2,4-diiodo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F

No.	Ex.	W	X	Y	Z
89		2,5-diiodo	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
90		2,4-diiodo	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
91		2-iodo,5-SCH <sub>3</sub>	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
92		2-iodo,5-ethyl	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
93		2-iodo,5-propyl	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
94		2-chloro,5-SCH <sub>3</sub>	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
95		2-chloro,5-ethyl	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
96		2-chloro,5-propyl	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
97		2,5-SCH <sub>3</sub>	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
98		2-iodo, 4-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
99		2-iodo, 3-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
100		2-iodo, 6-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
101		2-chloro, 3-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
102		2-chloro, 4-fluoro, 5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
103		2-chloro-3,4,5-trimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
104		2,3-diiodo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
105		2,5-dichloro	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
106		2,5-dibromo	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
107		2-iodo,4-chloro,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
108		2-iodo,4-bromo,5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F

No.	Ex.	W	X	Y	Z
109		2,5-dimethoxy	S	pent-4-ynyl	H
110		2,3,5-trimethoxy	S	pent-4-ynyl	H
112		3,4,5-trimethoxy	S	pent-4-ynyl	H
113	9.7	2-iodo,5-methoxy	S	pent-4-ynyl	H
114		2-bromo,5-methoxy	S	pent-4-ynyl	H
115		2-chloro,5-methoxy	S	pent-4-ynyl	H
116		2,4-diiodo,5-methoxy	S	pent-4-ynyl	H
117		2,5-diiodo	S	pent-4-ynyl	H
118		2,4-diiodo	S	pent-4-ynyl	H
119		2-iodo,5-SCH <sub>3</sub>	S	pent-4-ynyl	H
120		2-iodo,5-ethyl	S	pent-4-ynyl	H
121		2-iodo,5-propyl	S	pent-4-ynyl	H
122		2-chloro,5-SCH <sub>3</sub>	S	pent-4-ynyl	H
123		2-chloro,5-ethyl	S	pent-4-ynyl	H
124		2-chloro,5-propyl	S	pent-4-ynyl	H
125		2,5-SCH <sub>3</sub>	S	pent-4-ynyl	H
126		2-iodo, 4-fluoro, 5-methoxy	S	pent-4-ynyl	H
127		2-iodo, 3-fluoro, 5-methoxy	S	pent-4-ynyl	H
128		2-iodo, 6-fluoro, 5-methoxy	S	pent-4-ynyl	H
129		2-chloro, 3-fluoro, 5-methoxy	S	pent-4-ynyl	H
130		2-chloro, 4-fluoro, 5-methoxy	S	pent-4-ynyl	H

No.	Ex.	W	X	Y	Z
131		2-chloro-3,4,5-trimethoxy	S	pent-4-ynyl	H
132		2,3-diiodo,5-methoxy	S	pent-4-ynyl	H
133		2,5-dichloro	S	pent-4-ynyl	H
134		2,5-dibromo	S	pent-4-ynyl	H
135		2-iodo,4-chloro,5-methoxy	S	pent-4-ynyl	H
136		2-iodo,4-bromo,5-methoxy	S	pent-4-ynyl	H
137		2,5-dimethoxy	S	pent-4-ynyl	F
138		2,3,5-trimethoxy	S	pent-4-ynyl	F
139		3,4,5-trimethoxy	S	pent-4-ynyl	F
140		2-iodo,5-methoxy	S	pent-4-ynyl	F
141		2-bromo,5-methoxy	S	pent-4-ynyl	F
142		2-chloro,5-methoxy	S	pent-4-ynyl	F
143		2,4-diiodo,5-methoxy	S	pent-4-ynyl	F
144		2,5-diiodo	S	pent-4-ynyl	F
145		2,4-diiodo	S	pent-4-ynyl	F
146		2-iodo,5-SCH <sub>3</sub>	S	pent-4-ynyl	F
147		2-iodo,5-ethyl	S	pent-4-ynyl	F
148		2-iodo,5-propyl	S	pent-4-ynyl	F
149		2-chloro,5-SCH <sub>3</sub>	S	pent-4-ynyl	F
150		2-chloro,5-ethyl	S	pent-4-ynyl	F
151		2-chloro,5-propyl	S	pent-4-ynyl	F
152		2,5-SCH <sub>3</sub>	S	pent-4-ynyl	F
153		2-iodo, 4-fluoro, 5-	S	pent-4-ynyl	F

No.	Ex.	W	X	Y	Z
		methoxy			
154		2-iodo, 3-fluoro, 5-methoxy	S	pent-4-ynyl	F
155		2-iodo, 6-fluoro, 5-methoxy	S	pent-4-ynyl	F
156		2-chloro, 3-fluoro, 5-methoxy	S	pent-4-ynyl	F
157		2-chloro, 4-fluoro, 5-methoxy	S	pent-4-ynyl	F
158		2,3,4,5-tetraiodo	S	pent-4-ynyl	F
159		2,3-diiodo,5-methoxy	S	pent-4-ynyl	F
160		2,5-dichloro	S	pent-4-ynyl	F
161		2,5-dibromo	S	pent-4-ynyl	F
162		2-iodo,4-chloro,5-methoxy	S	pent-4-ynyl	F
163		2-iodo,4-bromo,5-methoxy	S	pent-4-ynyl	F
164		2,5-dimethoxy	S	4-methyl-pent-3-enyl	H
165		2,3,5-trimethoxy	S	4-methyl-pent-3-enyl	H
166		3,4,5-trimethoxy	S	4-methyl-pent-3-enyl	H
167	9.6	2-iodo,5-methoxy	S	4-methyl-pent-3-enyl	H
168		2-bromo,5-methoxy	S	4-methyl-pent-3-enyl	H
169		2-chloro,5-methoxy	S	4-methyl-pent-3-enyl	H
170		2,4-diiodo,5-methoxy	S	4-methyl-pent-3-enyl	H
171		2,5-diiodo	S	4-methyl-pent-3-enyl	H
172		2,4-diiodo	S	4-methyl-pent-3-enyl	H
173		2-iodo,5-SCH <sub>3</sub>	S	4-methyl-pent-3-enyl	H

No.	Ex.	W	X	Y	Z
174		2-iodo,5-ethyl	S	4-methyl-pent-3-enyl	H
175		2-iodo,5-propyl	S	4-methyl-pent-3-enyl	H
176		2-chloro,5-SCH <sub>3</sub>	S	4-methyl-pent-3-enyl	H
177		2-chloro,5-ethyl	S	4-methyl-pent-3-enyl	H
178		2-chloro,5-propyl	S	4-methyl-pent-3-enyl	H
179		2,5-SCH <sub>3</sub>	S	4-methyl-pent-3-enyl	H
180		2-iodo, 4-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	H
181		2-iodo, 3-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	H
182		2-iodo, 6-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	H
183		2-chloro, 3-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	H
184		2-chloro, 4-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	H
185		2-chloro-3,4,5-trimethoxy	S	4-methyl-pent-3-enyl	H
186		2,3-diiodo,5-methoxy	S	4-methyl-pent-3-enyl	H
187		2,5-dichloro	S	4-methyl-pent-3-enyl	H
188		2,5-dibromo	S	4-methyl-pent-3-enyl	H
189		2-iodo,4-chloro,5-methoxy	S	4-methyl-pent-3-enyl	H
190		2-iodo,4-bromo,5-methoxy	S	4-methyl-pent-3-enyl	H
191		2,5-dimethoxy	S	4-methyl-pent-3-enyl	F
192		2,3,5-trimethoxy	S	4-methyl-pent-3-enyl	F
193		3,4,5-trimethoxy	S	4-methyl-pent-3-enyl	F



No.	Ex.	W	X	Y	Z
194		2-iodo,5-methoxy	S	4-methyl-pent-3-enyl	F
195		2-bromo,5-methoxy	S	4-methyl-pent-3-enyl	F
196		2-chloro,5-methoxy	S	4-methyl-pent-3-enyl	F
197		2,4-diiodo,5-methoxy	S	4-methyl-pent-3-enyl	F
198		2,5-diiodo	S	4-methyl-pent-3-enyl	F
199		2,4-diiodo	S	4-methyl-pent-3-enyl	F
200		2-iodo,5-SCH <sub>3</sub>	S	4-methyl-pent-3-enyl	F
201		2-iodo,5-ethyl	S	4-methyl-pent-3-enyl	F
202		2-iodo,5-propyl	S	4-methyl-pent-3-enyl	F
203		2-chloro,5-SCH <sub>3</sub>	S	4-methyl-pent-3-enyl	F
204		2-chloro,5-ethyl	S	4-methyl-pent-3-enyl	F
205		2-chloro,5-propyl	S	4-methyl-pent-3-enyl	F
206		2,5-SCH <sub>3</sub>	S	4-methyl-pent-3-enyl	F
207		2-iodo, 4-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	F
208		2-iodo, 3-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	F
209		2-iodo, 6-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	F
210		2-chloro, 3-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	F
211		2-chloro, 4-fluoro, 5-methoxy	S	4-methyl-pent-3-enyl	F
212		2,3,4,5-tetraiodo	S	4-methyl-pent-3-enyl	F
213		2,3-diiodo,5-methoxy	S	4-methyl-pent-3-enyl	F
214		2,5-dichloro	S	4-methyl-pent-3-enyl	F

No.	Ex.	W	X	Y	Z
215		2,5-dibromo	S	4-methyl-pent-3-enyl	F
216		2-Iodo,4-chloro,5-methoxy	S	4-methyl-pent-3-enyl	F
217		2-iodo,4-bromo,5-methoxy	S	4-methyl-pent-3-enyl	F
218	11.2	2,5-dimethoxy	S	Pentyl	H
219	9.2	2,5-dimethoxy	S	Butyl	H
220	11.1	2,5-dimethoxy	S	H	H
221	11.3	2,5-dimethoxy	S	pent-4-ynyl	H
222	11.4	2,5-dimethoxy	S	butyronitrile	H
223	11.5	2,5-dimethoxy	S	3,3,3-trifluoropropyl	H
224	11.6	2,5-dimethoxy	S	4-chlorobutyl	H
225	11.7	2,5-dimethoxy	S	4-acetoxybutyl	H
226	11.8	2,5-dimethoxy	S	5-bromopentyl	H
227	11.9	2,5-dimethoxy	S	2-[1,3]dioxolan-2-yl-ethyl	H
228	11.10	2,5-dimethoxy	S	4-methyl-pent-3-enyl	H
229	11.11	2,5-dimethoxy	S	4-pentene	H
230	11.12	2,5-dimethoxy	S	3-hydroxypropyl	H
231	10.1	2,5-dimethoxy	S	4-methyl-pent-3-enyl	NH <sub>2</sub>
232	10.2	2,5-dimethoxy	S	4-methyl-pent-3-enyl	F
233	9.1	3-hydroxy	S	butyl	H
234	9.3	3-methoxy	S	butyl	H
235	9.4	2-iodo-5-methoxy	S	butyl	H
236	9.5	4-iodo-5-methoxy	S	Butyl	H
237		2,5-dimethoxy	CH <sub>2</sub>	Pentyl	H

No.	Ex.	W	X	Y	Z
238	1.1	2,5-dimethoxy	CH <sub>2</sub>	Butyl	H
239	3.8	2,5-dimethoxy	CH <sub>2</sub>	4-ethylaminobutyl	H
240		2,5-dimethoxy	CH <sub>2</sub>	pent-4-ynyl	H
241		2,5-dimethoxy	CH <sub>2</sub>	butyronitrile	H
242		2,5-dimethoxy	CH <sub>2</sub>	3,3,3-trifluoropropyl	H
243	3.1	2,5-dimethoxy	CH <sub>2</sub>	4-chlorobutyl	H
244	3.11	2,5-dimethoxy	CH <sub>2</sub>	5-acetoxypentyl	H
245	3.5	2,5-dimethoxy	CH <sub>2</sub>	5-bromopentyl	H
246	3.3	2,5-dimethoxy	CH <sub>2</sub>	2-[1,3]dioxolan-2-yl-ethyl	H
247	3.12	2,5-dimethoxy	CH <sub>2</sub>	3,3,3-trifluoropropyl	H
248	3.13	2,5-dimethoxy	CH <sub>2</sub>	4-pentene	H
249		2,5-dimethoxy	CH <sub>2</sub>	3-hydroxypropyl	H
250		2,5-dimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	NH <sub>2</sub>
251	2.2	2,5-dimethoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
252		2,5-dimethoxy	CH <sub>2</sub>	hexyl	H
253		2,5-dimethoxy	CH <sub>2</sub>	heptyl	H
254		2,5-dimethoxy	CH <sub>2</sub>	3-cyclopropylpropyl	H
255		2,5-dimethoxy	CH <sub>2</sub>	3-N,N-dimethylpropyl	H
256		2,5-dimethoxy	CH <sub>2</sub>	pentyl	F
257		2,5-dimethoxy	CH <sub>2</sub>	butyl	F
258		2,5-dimethoxy	CH <sub>2</sub>	4-ethylaminobutyl	F
259	2.1	2,5-dimethoxy	CH <sub>2</sub>	pent-4-ynyl	F
260		2,5-dimethoxy	CH <sub>2</sub>	butyronitrile	F
261		2,5-dimethoxy	CH <sub>2</sub>	3,3,3-trifluoropropyl	F

No.	Ex.	W	X	Y	Z
262		2,5-dimethoxy	CH <sub>2</sub>	4-chlorobutyl	F
263		2,5-dimethoxy	CH <sub>2</sub>	4-acetoxybutyl	F
264	2.4	2,5-dimethoxy	CH <sub>2</sub>	5-bromopentyl	F
265		2,5-dimethoxy	CH <sub>2</sub>	2-[1,3]dioxolan-2-yl-ethyl	F
266		2,5-dimethoxy	CH <sub>2</sub>	3,3,3-trifluoropropyl	F
267		2,5-dimethoxy	CH <sub>2</sub>	4-pentene	F
268		2,5-dimethoxy	CH <sub>2</sub>	3-hydroxypropyl	F
269		2,5-dimethoxy	CH <sub>2</sub>	4-cyclopropylbutyl	F
270		2,5-dimethoxy	CH <sub>2</sub>	4-ethyl-pent-3-enyl	F
271		2,5-dimethoxy	CH <sub>2</sub>	hexyl	F
272		2,5-dimethoxy	CH <sub>2</sub>	heptyl	F
273		2,5-dimethoxy	CH <sub>2</sub>	3-cyclopropylpropyl	F
274		2,5-dimethoxy	CH <sub>2</sub>	3-N,N-dimethylpropyl	F
275		2,5-dimethoxy	O	pentyl	F
276		2,5-dimethoxy	O	butyl	F
277		2,5-dimethoxy	O	H	F
278		2,5-dimethoxy	O	pent-4-ynyl	F
279		2,5-dimethoxy	O	butyronitrile	F
280		2,5-dimethoxy	O	3,3,3-trifluoropropyl	F
281		2,5-dimethoxy	O	4-chlorobutyl	F
282		2,5-dimethoxy	O	4-acetoxybutyl	F
283		2,5-dimethoxy	O	5-bromopentyl	F
284		2,5-dimethoxy	O	2-[1,3]dioxolan-2-yl-ethyl	F
285		2,5-dimethoxy	O	4-methyl-pent-3-enyl	F

No.	Ex.	W	X	Y	Z
286		2,5-dimethoxy	O	4-pentene	F
287		2,5-dimethoxy	O	3-hydroxypropyl	F
288		2,5-dimethoxy	O	4-methyl-pent-3-enyl	F
289		2,5-dimethoxy	O	4-ethyl-pent-3-enyl	F
290		2,5-dimethoxy	O	hexyl	F
291		2,5-dimethoxy	O	heptyl	F
292		2,5-dimethoxy	O	3-cyclopropylpropyl	F
293		2,5-dimethoxy	O	3-N,N-dimethylpropyl	F
294		2,5-dimethoxy	O	pentyl	H
295		2,5-dimethoxy	O	butyl	H
296		2,5-dimethoxy	O	H	H
297		2,5-dimethoxy	O	pent-4-ynyl	H
298		2,5-dimethoxy	O	butyronitrile	H
299		2,5-dimethoxy	O	3,3,3-trifluoropropyl	H
300		2,5-dimethoxy	O	4-chlorobutyl	H
301		2,5-dimethoxy	O	4-acetoxybutyl	H
302		2,5-dimethoxy	O	5-bromopentyl	H
303		2,5-dimethoxy	O	2-[1,3]dioxolan-2-yl-ethyl	H
304		2,5-dimethoxy	O	4-methyl-pent-3-enyl	H
305		2,5-dimethoxy	O	4-pentene	H
306		2,5-dimethoxy	O	3-hydroxypropyl	H
307		2,5-dimethoxy	O	4-cyclopropylbutyl	H
308		2,5-dimethoxy	O	4-ethyl-pent-3-enyl	H
309		2,5-dimethoxy	O	hexyl	H

No.	Ex.	W	X	Y	Z
310		2,5-dimethoxy	O	heptyl	H
311		2,5-dimethoxy	O	3-cyclopropylpropyl	H
312		2,5-dimethoxy	O	3-N,N-dimethylpropyl	H
313		2-iodo-5-methoxy	O	pentyl	F
314		2-iodo-5-methoxy	O	butyl	F
315		2-iodo-5-methoxy	O	H	F
316		2-iodo-5-methoxy	O	pent-4-ynyl	F
317		2-iodo-5-methoxy	O	butyronitrile	F
318		2-iodo-5-methoxy	O	3,3,3-trifluoropropyl	F
319		2-iodo-5-methoxy	O	4-chlorobutyl	F
320		2-iodo-5-methoxy	O	4-acetoxybutyl	F
321		2-iodo-5-methoxy	O	5-bromopentyl	F
322		2-iodo-5-methoxy	O	2-[1,3]dioxolan-2-yl-ethyl	F
323		2-iodo-5-methoxy	O	4-methyl-pent-3-enyl	F
324		2-iodo-5-methoxy	O	4-pentene	F
325		2-iodo-5-methoxy	O	3-hydroxypropyl	F
326		2-iodo-5-methoxy	O	4-cyclopropylbutyl	F
327		2-iodo-5-methoxy	O	4-ethyl-pent-3-enyl	F
328		2-iodo-5-methoxy	O	hexyl	F
329		2-iodo-5-methoxy	O	heptyl	F
330		2-iodo-5-methoxy	O	3-cyclopropylpropyl	F
331		2-iodo-5-methoxy	O	3-N,N-dimethylpropyl	F
332		2-iodo-5-methoxy	O	pentyl	H
333		2-iodo-5-methoxy	O	butyl	H

No.	Ex.	W	X	Y	Z
334		2-iodo-5-methoxy	O	H	H
335		2-iodo-5-methoxy	O	pent-4-ynyl	H
336		2-iodo-5-methoxy	O	butyronitrile	H
337		2-iodo-5-methoxy	O	3,3,3-trifluoropropyl	H
338		2-iodo-5-methoxy	O	4-chlorobutyl	H
339		2-iodo-5-methoxy	O	4-acetoxybutyl	H
340		2-iodo-5-methoxy	O	5-bromopentyl	H
341		2-iodo-5-methoxy	O	2-[1,3]dioxolan-2-yl-ethyl	H
342		2-iodo-5-methoxy	O	4-methyl-pent-3-enyl	H
343		2-iodo-5-methoxy	O	4-pentene	H
344		2-iodo-5-methoxy	O	3-hydroxypropyl	H
345		2-iodo-5-methoxy	O	4-cyclopropylbutyl	H
346		2-iodo-5-methoxy	O	4-ethyl-pent-3-enyl	H
347		2-iodo-5-methoxy	O	hexyl	H
348		2-iodo-5-methoxy	O	heptyl	H
349		2-iodo-5-methoxy	O	3-cyclopropylpropyl	H
350		2-iodo-5-methoxy	O	3-N,N-dimethylpropyl	H
351		2-iodo-5-methoxy	NH	pentyl	F
352		2-iodo-5-methoxy	NH	butyl	F
353		2-iodo-5-methoxy	NH	H	F
354		2-iodo-5-methoxy	NH	pent-4-ynyl	F
355		2-iodo-5-methoxy	NH	butyronitrile	F
356		2-iodo-5-methoxy	NH	3,3,3-trifluoropropyl	F
357		2-iodo-5-methoxy	NH	4-chlorobutyl	F

No.	Ex.	W	X	Y	Z
358		2-iodo-5-methoxy	NH	4-acetoxybutyl	F
359		2-iodo-5-methoxy	NH	5-bromopentyl	F
360		2-iodo-5-methoxy	NH	2-[1,3]dioxolan-2-yl-ethyl	F
361		2-iodo-5-methoxy	NH	4-methyl-pent-3-enyl	F
362		2-iodo-5-methoxy	NH	4-pentene	F
363		2-iodo-5-methoxy	NH	3-hydroxypropyl	F
364		2-iodo-5-methoxy	NH	4-cyclopropylbutyl	F
365		2-iodo-5-methoxy	NH	4-ethyl-pent-3-enyl	F
366		2-iodo-5-methoxy	NH	hexyl	F
367		2-iodo-5-methoxy	NH	heptyl	F
368		2-iodo-5-methoxy	NH	3-cyclopropylpropyl	F
369		2-iodo-5-methoxy	NH	3-N,N-dimethylpropyl	F
370		2-iodo-5-methoxy	NH	pentyl	H
371		2-iodo-5-methoxy	NH	butyl	H
372		2-iodo-5-methoxy	NH	H	H
373		2-iodo-5-methoxy	NH	pent-4-ynyl	H
374		2-iodo-5-methoxy	NH	butyronitrile	H
375		2-iodo-5-methoxy	NH	3,3,3-trifluoropropyl	H
376		2-iodo-5-methoxy	NH	4-chlorobutyl	H
377		2-iodo-5-methoxy	NH	4-acetoxybutyl	H
378		2-iodo-5-methoxy	NH	5-bromopentyl	H
379		2-iodo-5-methoxy	NH	2-[1,3]dioxolan-2-yl-ethyl	H
380		2-iodo-5-methoxy	NH	4-methyl-pent-3-enyl	H
381		2-iodo-5-methoxy	NH	4-pentene	H



No.	Ex.	W	X	Y	Z
382		2-iodo-5-methoxy	NH	3-hydroxypropyl	H
383		2-iodo-5-methoxy	NH	4-cyclopropylbutyl	H
384		2-iodo-5-methoxy	NH	4-ethyl-pent-3-enyl	H
385		2-iodo-5-methoxy	NH	hexyl	H
386		2-iodo-5-methoxy	NH	heptyl	H
387		2-iodo-5-methoxy	NH	3-cyclopropylpropyl	H
388		2-iodo-5-methoxy	NH	3-N,N-dimethylpropyl	H
389		2,5-dimethoxy	NH	pentyl	F
390		2,5-dimethoxy	NH	butyl	F
391		2,5-dimethoxy	NH	H	F
392		2,5-dimethoxy	NH	pent-4-ynyl	F
393		2,5-dimethoxy	NH	butyronitrile	F
394		2,5-dimethoxy	NH	3,3,3-trifluoropropyl	F
395		2,5-dimethoxy	NH	4-chlorobutyl	F
396		2,5-dimethoxy	NH	4-acetoxybutyl	F
397		2,5-dimethoxy	NH	5-bromopentyl	F
398		2,5-dimethoxy	NH	2-[1,3]dioxolan-2-yl-ethyl	F
399		2,5-dimethoxy	NH	4-methyl-pent-3-enyl	F
400		2,5-dimethoxy	NH	4-pentene	F
401		2,5-dimethoxy	NH	3-hydroxypropyl	F
402		2,5-dimethoxy	NH	4-cyclopropylbutyl	F
403		2,5-dimethoxy	NH	4-ethyl-pent-3-enyl	F
404		2,5-dimethoxy	NH	hexyl	F
405		2,5-dimethoxy	NH	heptyl	F

No.	Ex.	W	X	Y	Z
406		2,5-dimethoxy	NH	3-cyclopropylpropyl	F
407		2,5-dimethoxy	NH	3-N,N-dimethylpropyl	F
408		2,5-dimethoxy	NH	pentyl	H
409		2,5-dimethoxy	NH	butyl	H
410		2,5-dimethoxy	NH	H	H
411		2,5-dimethoxy	NH	pent-4-ynyl	H
412		2,5-dimethoxy	NH	butyronitrile	H
413		2,5-dimethoxy	NH	3,3,3-trifluoropropyl	H
414		2,5-dimethoxy	NH	4-chlorobutyl	H
415		2,5-dimethoxy	NH	4-acetoxybutyl	H
416		2,5-dimethoxy	NH	5-bromopentyl	H
417		2,5-dimethoxy	NH	2-[1,3]dioxolan-2-yl-ethyl	H
418		2,5-dimethoxy	NH	4-methyl-pent-3-enyl	H
419		2,5-dimethoxy	NH	4-pentene	H
420		2,5-dimethoxy	NH	3-hydroxypropyl	H
421		2,5-dimethoxy	NH	4-cyclopropylbutyl	H
422		2,5-dimethoxy	NH	4-ethyl-pent-3-enyl	H
423		2,5-dimethoxy	NH	hexyl	H
424		2,5-dimethoxy	NH	heptyl	H
425		2,5-dimethoxy	NH	3-cyclopropylpropyl	H
426		2,5-dimethoxy	NH	3-N,N-dimethylpropyl	H
427		2,5-dimethoxy	S	pentyl	F
428		2,5-dimethoxy	S	4-ethylaminobutyl	H
429		2,5-dimethoxy	S	4-ethylaminobutyl	F

No.	Ex.	W	X	Y	Z
430		2,5-dimethoxy	S	pent-4-ynyl	F
431		2,5-dimethoxy	S	butyronitrile	F
432		2,5-dimethoxy	S	3,3,3-trifluoropropyl	F
433		2,5-dimethoxy	S	4-chlorobutyl	F
434		2,5-dimethoxy	S	4-acetoxybutyl	F
435		2,5-dimethoxy	S	5-bromopentyl	F
436		2,5-dimethoxy	S	2-[1,3]dioxolan-2-yl-ethyl	F
437		2,5-dimethoxy	S	butyl	F
438		2,5-dimethoxy	S	4-pentene	F
439		2,5-dimethoxy	S	3-hydroxypropyl	F
440		2,5-dimethoxy	S	4-methyl-pent-3-enyl	F
441		2,5-dimethoxy	S	4-ethyl-pent-3-enyl	F
442		3-hydroxy	S	butyl	F
443		2,5-dimethoxy	S	2-(dimethyl-bicyclo[3.1.1]hept-2-en-2-yl)-ethyl	F
444		2-Iodo-5-methoxy	S	butyl	F
445		4-Iodo-5-methoxy	S	butyl	F
446		2,5-dimethoxy	S	hexan-6-ol	H
447		2,5-dimethoxy	S	pentan-5-ol	H
448		2,5-dimethoxy	S	butan-4-ol	H
449		2,5-dimethoxy	S	hexan-6-ol	F
450		2,5-dimethoxy	S	pentan-5-ol	F
451		2,5-dimethoxy	S	butan-4-ol	F
452		2-iodo-5-methoxy	S	hexan-6-ol	H

No.	Ex.	W	X	Y	Z
453		2-iodo-5-methoxy	S	pentan-5-ol	H
454		2-iodo-5-methoxy	S	butan-4-ol	H
455		2-iodo-5-methoxy	S	hexan-6-ol	F
456		2-iodo-5-methoxy	S	pentan-5-ol	F
457		2-iodo-5-methoxy	S	butan-4-ol	F
458		2-iodo-5-methoxy	S	hexan-6-ol	H
459		2-iodo-5-methoxy	S	pentan-5-ol	H
460		2,5-dimethoxy	CH <sub>2</sub>	hexan-6-ol	H
461		2,5-dimethoxy	CH <sub>2</sub>	pentan-5-ol	H
462		2,5-dimethoxy	CH <sub>2</sub>	butan-4-ol	H
463	3.9	2,5-dimethoxy	CH <sub>2</sub>	hexan-6-ol	F
464		2,5-dimethoxy	CH <sub>2</sub>	pentan-5-ol	F
465		2,5-dimethoxy	CH <sub>2</sub>	butan-4-ol	F
466		2-iodo-5-methoxy	CH <sub>2</sub>	hexan-6-ol	H
467		2-iodo-5-methoxy	CH <sub>2</sub>	pentan-5-ol	H
468		2-iodo-5-methoxy	CH <sub>2</sub>	butan-4-ol	H
469		2-iodo-5-methoxy	CH <sub>2</sub>	hexan-6-ol	F
470		2-iodo-5-methoxy	CH <sub>2</sub>	pentan-5-ol	F
471		2-iodo-5-methoxy	CH <sub>2</sub>	butan-4-ol	F
472		2-iodo-5-methoxy	CH <sub>2</sub>	hexan-6-ol	H
473		2-iodo-5-methoxy	CH <sub>2</sub>	pentan-5-ol	H
474	2.3	2,5-dimethoxy	CH <sub>2</sub>	pent-4-enyl	F
475	3.6	2,5-dimethoxy	CH <sub>2</sub>	5-bromo-3-methyl-pentyl	H
476	3.6	2,5-dimethoxy	CH <sub>2</sub>	5-chloro-pentyl	H

No.	Ex.	W	X	Y	Z
477	3.10	2,5-dimethoxy	CH <sub>2</sub>	dimethyl-bicyclo[3.1.1]hept-2-en-2-yl)-ethyl	H
478		2-iodo-5-methoxy	CH <sub>2</sub>	pentyl	H
479	4.1	2-iodo-5-methoxy	CH <sub>2</sub>	butyl	H
480		2-iodo-5-methoxy	CH <sub>2</sub>	4-ethylaminobutyl	H
481		2-iodo-5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
482		2-iodo-5-methoxy	CH <sub>2</sub>	butyronitrile	H
483		2-iodo-5-methoxy	CH <sub>2</sub>	3,3,3-trifluoropropyl	H
484		2-iodo-5-methoxy	CH <sub>2</sub>	4-chlorobutyl	H
485		2-iodo-5-methoxy	CH <sub>2</sub>	5-acetoxypentyl	H
486		2-iodo-5-methoxy	CH <sub>2</sub>	5-bromopentyl	H
487		2-iodo-5-methoxy	CH <sub>2</sub>	2-[1,3]dioxolan-2-yl-ethyl	H
488		2-iodo-5-methoxy	CH <sub>2</sub>	3,3,3-trifluoropropyl	H
489		2-iodo-5-methoxy	CH <sub>2</sub>	4-pentene	H
490		2-iodo-5-methoxy	CH <sub>2</sub>	3-hydroxypropyl	H
491	4.2	5-iodo-2-methoxy	CH <sub>2</sub>	butyl	H
492		2-iodo-5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	F
493		2-iodo-5-methoxy	CH <sub>2</sub>	hexyl	H
494		2-iodo-5-methoxy	CH <sub>2</sub>	heptyl	H
495		2-iodo-5-methoxy	CH <sub>2</sub>	3-cyclopropylpropyl	H
496		2-iodo-5-methoxy	CH <sub>2</sub>	3-N,N-dimethylpropyl	H
497	4.3	5-ethyl-2-methoxy	CH <sub>2</sub>	butyl	H
498	4.4	2-bromo-5-methoxy	CH <sub>2</sub>	butyl	H
499	4.5	2-bromo-5-methoxy	CH <sub>2</sub>	4-methyl-pent-3-enyl	H

No.	Ex.	W	X	Y	Z
500	4.6	2-bromo-5-methoxy	CH <sub>2</sub>	pent-4-ynyl	H
501	5.1	5-ethyl-2-methoxy	CH <sub>2</sub>	butyl	H
502	5.2	5-butyl-2-methoxy	CH <sub>2</sub>	butyl	H
503	5.3	5-vinyl-2-methoxy	CH <sub>2</sub>	butyl	H
504	6.1	2,5-dimethoxy-4-nitro	CH <sub>2</sub>	pent-4-ynyl	F
505	6.2	3,5-dimethoxy-2-nitro	CH <sub>2</sub>	butyl	H
506	6.3	3,5-dimethoxy-4-amino	CH <sub>2</sub>	4-methyl-pent-3-enyl	H
507	6.4	2,5-dimethoxy-4-amino	CH <sub>2</sub>	butyl	H
508	6.5	3,5-dimethoxy-2-amino	CH <sub>2</sub>	butyl	H
509	6.6	4-methoxy benzaldehyde-O- methyl-oxime	CH <sub>2</sub>	butyl	H
510	7.1	4-methoxy benzaldehyde	CH <sub>2</sub>	butyl	H
511	8.1	3,4-dichloro benzyl	CH <sub>2</sub>	butyl	H
512		2-iodo-5-methoxy	S	pentyl	H
513		2-iodo-5-methoxy	S	butyl	H
514		2-iodo-5-methoxy	S	4-ethylaminobutyl	H
515		2-iodo-5-methoxy	S	pent-4-ynyl	H
516		2-iodo-5-methoxy	S	butyronitrile	H
517		2-iodo-5-methoxy	S	3,3,3-trifluoropropyl	H

No.	Ex.	W	X	Y	Z
518		2-iodo-5-methoxy	S	4-chlorobutyl	H
519		2-iodo-5-methoxy	S	5-acetoxypentyl	H
520		2-iodo-5-methoxy	S	5-bromopentyl	H
521		2-iodo-5-methoxy	S	2-[1,3]dioxolan-2-yl-ethyl	H
522		2-iodo-5-methoxy	S	3,3,3-trifluoropropyl	H
523		2-iodo-5-methoxy	S	4-pentene	H
524		2-iodo-5-methoxy	S	3-hydroxypropyl	H
525		2-iodo-5-methoxy	S	4-methyl-pent-3-enyl	NH <sub>2</sub>
526		2-iodo-5-methoxy	S	4-methyl-pent-3-enyl	F
527		2-iodo-5-methoxy	S	hexyl	H
528		2-iodo-5-methoxy	S	heptyl	H
529		2-iodo-5-methoxy	S	3-cyclopropylpropyl	H
530		2-iodo-5-methoxy	S	3-N,N-dimethylpropyl	H
531		2-iodo-5-methoxy	S	pentyl	F
532		2-iodo-5-methoxy	S	butyl	F
533		2-iodo-5-methoxy	S	4-ethylaminobutyl	F
534		2-iodo-5-methoxy	S	pent-4-ynyl	F
536		2-iodo-5-methoxy	S	butyronitrile	F
537		2-iodo-5-methoxy	S	3,3,3-trifluoropropyl	F
538		2-iodo-5-methoxy	S	4-chlorobutyl	F
539		2-iodo-5-methoxy	S	4-acetoxybutyl	F
540		2-iodo-5-methoxy	S	5-bromopentyl	F
541		2-iodo-5-methoxy	S	2-[1,3]dioxolan-2-yl-ethyl	F
542		2-iodo-5-methoxy	S	3,3,3-trifluoropropyl	F

No.	Ex.	W	X	Y	Z
543		2-iodo-5-methoxy	S	4-pentene	F
544		2-iodo-5-methoxy	S	3-hydroxypropyl	F
545		2-iodo-5-methoxy	S	4-cyclopropylbutyl	F
546		2-iodo-5-methoxy	S	4-ethyl-pent-3-enyl	F
547		2-iodo-5-methoxy	S	hexyl	F
548		2-iodo-5-methoxy	S	heptyl	F
549		2-iodo-5-methoxy	S	3-cyclopropylpropyl	F
550		2-iodo-5-methoxy	S	3-N,N-dimethylpropyl	F
551		2-iodo-5-methoxy	S	pentyl	Cl
552		2-iodo-5-methoxy	S	butyl	Cl
553		2-iodo-5-methoxy	S	4-ethylaminobutyl	Cl
554		2-iodo-5-methoxy	S	pent-4-ynyl	Cl
555		2-iodo-5-methoxy	S	butyronitrile	Cl
556		2-iodo-5-methoxy	S	3,3,3-trifluoropropyl	Cl
557		2-iodo-5-methoxy	S	4-chlorobutyl	Cl
558		2-iodo-5-methoxy	S	4-acetoxybutyl	Cl
559		2-iodo-5-methoxy	S	5-bromopentyl	Cl
560		2-iodo-5-methoxy	S	2-[1,3]dioxolan-2-yl-ethyl	Cl
561		2-iodo-5-methoxy	S	3,3,3-trifluoropropyl	Cl
562		2-iodo-5-methoxy	S	4-pentene	Cl
563		2-iodo-5-methoxy	S	3-hydroxypropyl	Cl
564		2-iodo-5-methoxy	S	4-cyclopropylbutyl	Cl
565		2-iodo-5-methoxy	S	4-ethyl-pent-3-enyl	Cl
566		2-iodo-5-methoxy	S	hexyl	Cl



No.	Ex.	W	X	Y	Z
567		2-iodo-5-methoxy	S	heptyl	Cl
568		2-iodo-5-methoxy	S	3-cyclopropylpropyl	Cl
569		2-iodo-5-methoxy	S	3-N,N-dimethylpropyl	Cl
570		2-iodo-5-methoxy	S	4-methyl-pent-3-enyl	NH <sub>2</sub>
571		2-iodo-5-methoxy	S	4-methyl-pent-3-enyl	Cl
572		2-iodo-5-methoxy	S	pent-4-ynyl	Cl
573	9.11	2,5-dichloro	S	butyl	H
574	9.12	2,4,5-trichloro	S	butyl	H
575	2.5	2,5-dimethoxy	CH <sub>2</sub>	pent-4-ynyl	Cl

Particularly preferred in Table 1 are compounds 1, 4, 31, 55, 58, 85, 113, 137, 140, 167, 191, 194, 221, 224, 226, 228, 232, 234, 235, 236, 239, 245, 248, 251, 259, 452, 453, 454, 455, 456, 457, 463, 469, 470, 471, 571 and 572, with the most preferred compounds being 1, 85, 113, 137, 140, 167, 191, 194, 221, 228, 239, 251, 259, 452, 453, 455, 571 and 572.

Other compounds of the invention are based on the following formula, having illustrative species as described in Table 2:





Table 2

No.	Ex.	E	X	L	Z
576	9.8	H	S	S	H
577	9.9	5-chloro	S	S	H
578	9.10	5-methoxy	S	S	H


The foregoing aspects and embodiments can also include tautomers of the compounds. For example, referring to structures I-IV, above, structure II is a tautomeric form of structure I when A is OH (OR<sup>3</sup> where R<sup>3</sup> is H), structure III is another tautomeric form of structure I (when Z is OH), and structure IV is yet another tautomeric form of structure I in the event that both A and Z are OH. All four tautomeric possibilities can be represented essentially as shown in structure I, except that dashed lines appear between atoms 1 and 6 and 2 and 3, and between A and 6 and Z and 2, e.g., as shown:



The foregoing aspects and embodiments can also include prodrugs and/or pharmaceutically acceptable salts of the compounds shown.

The foregoing aspects and embodiments can also include varying levels and types of substitution on one or more of entities A, B, Q, W, X, Y and Z, above, where that entity is not already solely hydrogen but rather is a multi-atom substituent that contains one or more hydrogens that can be substituted for, e.g., with a halogen or combination of other atoms or chemical group(s). In some embodiments, there may be a range of from 0 to 25 or more collective substitutions (for all of A, B, Q, W, X, Y and Z), and any range in

between. The substitutions may be made or included at any point in the synthesis of the final compounds, as appropriate, including, e.g., in the starting reagents or intermediates of the reaction scheme(s) used, or following the synthesis of one final product to convert it into another. The following embodiments are illustrative:



5

In another aspect, the invention features pharmaceutical compositions containing one or more of the compounds or pharmaceutically acceptable salts thereof described for the preceding aspects. These additionally include one or more pharmaceutically acceptable carriers and/or excipients.

Another aspect of the invention features methods of making the compounds of the preceding aspects. These are described in greater detail in the next section and the examples to follow. Related aspects of the invention embrace intermediates of/in these synthetic methods to the extent they are novel, either alone or standing in the context of the specific synthesis objective.

In yet another aspect, the invention features methods of inhibiting an HSP90 molecule with a compound according to any of the previous aspects and embodiments. HSP90 proteins are highly conserved in nature (see, e.g., NCBI accession #'s P07900 and XM 004515 (human  $\alpha$  and  $\beta$  HSP90, respectively), P11499 (mouse), AAB2369 (rat), P46633 (chinese hamster), JC1468 (chicken), AAF69019 (flesh fly), AAC21566 (zebrafish), AAD30275 (salmon), O02075 (pig), NP 015084 (yeast), and CAC29071 (frog). Grp94 and Trap-1 are related molecules falling within the definition of an HSP90- as used herein. There are thus many different HSP90s, all with expected similar effect and inhibition capabilities. The HSP90 inhibitors of the invention may be specifically directed against an HSP90 of the specific host patient or may be identified based on reactivity against an HSP90 homolog from a different species, or an HSP90 variant. The methods feature contacting a cell having an HSP90 with a pharmaceutically effective amount of a compound or pharmaceutical composition according to any one of the preceding aspects. The cell is preferably a mammalian cell, and more preferably a human cell, although any cell-type from any life-form that contains an HSP90, including non-mammalian lines, is contemplated for the invention. The method can be "in vitro", e.g., contacting a cell line in culture, or else can be "in vivo", e.g., contacting a cell inside a live organism. One type of in vivo administration is made "in situ", or directly to a specific cell or group of cells within an organism, e.g, intratumorally. "Ex vivo" procedures are also envisioned wherein the cells are first removed from a patient, treated by contacting them with the compounds or compositions of the invention, and then administered back to a patient or "the" patient. The compounds and compositions can be administered in a variety of ways, e.g., intravenously, parenterally, orally, buccally, intramuscularly, sublingually, topically, by aerosol, subcutaneously, intramuscularly, intraperitoneally, rectally, vaginally, intratumorally, or peritumorally.

In some aspects, the compounds or compositions are administered to treat or prevent a cancer, e.g., a breast cancer, melanoma, lung cancer, etc. In some embodiments, these compounds may be used in combination with or as an adjuvant/sensitizer for any chemotherapy regimen. Such regimens may include the use of other anti-cancer compounds, e.g., taxol, Herceptin, Gleevac, etc. The additions may be made simultaneously or sequentially and, if the latter, in any order.

In other aspects, the compounds or compositions are used for non-oncology applications, e.g., treating inflammation, infectious disease, autoimmune disease, and ischemia.

Any of the above described aspects and embodiments of the invention can be combined where practical. The individual methods prescribed do not preclude the utilization of other, unspecified steps, and those of ordinary skill in the art will appreciate that additional steps and compounds may also be usefully incorporated within the spirit of the invention.

Advantages of the invention depend on the specific aspect and embodiment and may include one or more of: ease of synthesis and/or formulation, solubility, and IC<sub>50</sub> relative to previously existing compounds in the same or different classes of HSP90 inhibitors. Other advantages, aspects, and embodiments will be apparent from the figures, the detailed description and claims to follow.

## BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows IC<sub>50</sub> values for compounds of Table 3, Example 3, as measured using Her-2 degradation studies.

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions

A “pharmaceutically acceptable salt” may be prepared for any compound of the invention having a functionality capable of forming a salt, for example an acid or base functionality. Pharmaceutically acceptable salts may be derived from organic or inorganic acids and bases.


Compounds of the invention that contain one or more basic functional groups, e.g., amino or alkylamino, are capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable organic and inorganic acids. These salts can be prepared in situ during the final isolation and purification of the compounds of the invention, or by  
5 separately reacting a purified compound of the invention in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed. Examples of suitable acids include hydrochloric, hydrobromic, sulfuric, nitric, perchloric, fumaric, maleic, phosphoric, glycolic, gluconic, lactic, salicylic, succinic, toluene-p-sulfonic, tartaric, acetic, citric, methanesulfonic, formic, benzoic, malonic, naphthalene-2-sulfonic,  
10 benzenesulfonic, 1,2 ethanesulfonic acid (edisylate), galactosyl-d-gluconic acid, and the like. Other acids, such as oxalic acid, while not themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of this invention and their pharmaceutically acceptable acid addition salts. See, e.g., Berge et al. "Pharmaceutical Salts", J. Pharm. Sci. 66:1-19 (1977).

15 Compounds of the present invention that contain one or more acidic functional groups are capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable bases. The term "pharmaceutically-acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These salts can likewise be prepared in situ during the final isolation  
20 and purification of the compounds, or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation, with ammonia, or with a pharmaceutically-acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum  
25 salts and the like. Illustrative examples of some of the bases that can be used include sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate, and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like. See, for example, Berge et al., supra.

30 "Prodrugs" are derivative compounds derivatized by the addition of a group that endows greater solubility to the compound desired to be delivered. Once in the body, the

prodrug is typically acted upon by an enzyme, e.g., an esterase, amidase, or phosphatase, to generate the active compound. Suitable positions for derivatization of the compounds of the invention to create “prodrugs” include but are not limited to the Y group, the phenyl ring of the purines, and the Q group. Those of ordinary skill in the art have the knowledge and means to accomplish this without undue experimentation.

“Tautomers” are compounds whose structures differ in arrangements of atoms, but which exist in equilibrium. By way of example, the structure shown below and designated T is in equilibrium with a second tautomeric form designated T'.



The predominance of one tautomer versus another is controlled by factors which include but are not limited to the nature of the solvent, temperature, pressure, the presence or absence of other molecules, and the nature of substituents on the molecule having tautomeric forms.

The term “alkyl,” alone or in combination, refers to an optionally substituted straight-chain, optionally substituted branched-chain, or optionally substituted cyclic alkyl radical having from 1 to about 30 carbons, more preferably 1 to 12 carbons. Examples of alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, tert-amyl, pentyl, hexyl, heptyl, octyl and the like. The term “cycloalkyl” embraces cyclic configurations, is subsumed within the definition of alkyl and specifically refers to a monocyclic, bicyclic, tricyclic, and higher multicyclic alkyl radicals wherein each cyclic moiety has from 3 to about 8 carbon atoms. Examples of cycloalkyl radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like. A “lower alkyl” is a shorter alkyl, e.g., one containing from 1 to about 6 carbon atoms.

The term “alkenyl,” alone or in combination, refers to an optionally substituted straight-chain, optionally substituted branched-chain, or optionally substituted cyclic alkenyl hydrocarbon radical having one or more carbon-carbon double-bonds and having

from 2 to about 30 carbon atoms, more preferably 2 to about 18 carbons. Examples of alkenyl radicals include ethenyl, propenyl, butenyl, 1,4-butadienyl and the like. The term can also embrace cyclic alkenyl structures. A “lower alkenyl” refers to an alkenyl having from 2 to about 6 carbons.

5           The term “alkynyl,” alone or in combination, refers to an optionally substituted straight-chain, optionally substituted branched-chain, or cyclic alkynyl hydrocarbon radical having one or more carbon-carbon triple-bonds and having from 2 to about 30 carbon atoms, more preferably 2 to about 12 carbon atoms. The term also includes optionally substituted straight-chain or optionally substituted branched-chain hydrocarbon  
10 radicals having one or more carbon-carbon triple bonds and having from 2 to about 6 carbon atoms as well as those having from 2 to about 4 carbon atoms. Examples of alkynyl radicals include ethynyl, propynyl, butynyl and the like.

          The terms heteroalkyl, heteroalkenyl and heteroalkynyl include optionally substituted alkyl, alkenyl and alkynyl structures, as described above, and which have one  
15 or more skeletal chain atoms selected from an atom other than carbon, *e.g.*, oxygen, nitrogen, sulfur, phosphorous or combinations thereof.

          The term “carbon chain” may embrace any alkyl, alkenyl, alkynyl, or heteroalkyl, heteroalkenyl, or heteroalkynyl group, and may be linear, cyclic, or any combination thereof. If part of a linker and that linker comprises one or more rings as part of the core  
20 backbone, for purposes of calculating chain length, the “chain” only includes those carbon atoms that compose the bottom or top of a given ring and not both, and where the top and bottom of the ring(s) are not equivalent in length, the shorter distance shall be used in determining chain length. If the chain contains heteroatoms as part of the backbone, those atoms are not calculated as part of the carbon chain length.

25           The term “alkoxy,” alone or in combination, refers to an alkyl ether radical, alkyl-O-, wherein the term alkyl is defined as above. Examples of alkoxy radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy and the like.



The term "aryloxy," alone or in combination, refers to an aryl ether radical wherein the term aryl is defined as below. Examples of aryloxy radicals include phenoxy, benzyloxy and the like.

5 The term "alkylthio," alone or in combination, refers to an alkyl thio radical, alkyl-S-, wherein the term alkyl is defined as above.

The term "arylthio," alone or in combination, refers to an aryl thio radical, aryl-S-, wherein the term aryl is defined as below.

The term "oxo" refers to =O.

10 The term "aryl," alone or in combination, refers to an optionally substituted aromatic ring system. The term aryl includes monocyclic aromatic rings, polyaromatic rings and polycyclic aromatic ring systems containing from six to about twenty carbon atoms. The term aryl also includes monocyclic aromatic rings, polyaromatic rings and polycyclic ring systems containing from 6 to about 12 carbon atoms, as well as those containing from 6 to about 10 carbon atoms. The polyaromatic and polycyclic aromatic  
15 rings systems may contain from two to four rings. Examples of aryl groups include, without limitation, phenyl, biphenyl, naphthyl and anthryl ring systems.

The term "heteroaryl" refers to optionally substituted aromatic ring systems containing from about five to about 20 skeletal ring atoms and having one or more heteroatoms such as, for example, oxygen, nitrogen, sulfur, and phosphorus. The term  
20 heteroaryl also includes optionally substituted aromatic ring systems having from 5 to about 12 skeletal ring atoms, as well as those having from 5 to about 10 skeletal ring atoms. The term heteroaryl may include five- or six-membered heterocyclic rings, polycyclic heteroaromatic ring systems and polyheteroaromatic ring systems where the ring system has two, three or four rings. The terms heterocyclic, polycyclic  
25 heteroaromatic and polyheteroaromatic include ring systems containing optionally substituted heteroaromatic rings having more than one heteroatom as described above (*e.g.*, a six membered ring with two nitrogens), including polyheterocyclic ring systems of from two to four rings. The term heteroaryl includes ring systems such as, for example, furanyl, benzofuranyl, chromenyl, pyridyl, pyrrolyl, indolyl, quinoliny, N-alkyl pyrrolyl,

pyridyl-N-oxide, pyrimidoyl, pyrazinyl, imidazolyl, pyrazolyl, oxazolyl, benzothiophenyl, purinyl, indolizinyl, thienyl and the like.


The term “heteroarylalkyl” refers to a C1-C4 alkyl group containing a heteroaryl group, each of which may be optionally substituted.

5        The term “heteroarylthio” refers to the group -S-heteroaryl.

The term “acyloxy” refers to the ester group  $-\text{OC}(\text{O})-\text{R}$ , where R is H, alkyl, alkenyl, alkynyl, aryl, or arylalkyl, wherein the alkyl, alkenyl, alkynyl and arylalkyl groups may be optionally substituted.

10       The term “carboxy esters” refers to  $-\text{C}(\text{O})\text{OR}$  where R is alkyl, aryl or arylalkyl, wherein the alkyl, aryl and arylalkyl groups may be optionally substituted.

The term “carboxamido” refers to



15       where each of R and R' are independently selected from the group consisting of H, alkyl, aryl and arylalkyl, wherein the alkyl, aryl and arylalkyl groups may be optionally substituted.

The term “arylalkyl,” alone or in combination, refers to an alkyl radical as defined above in which one H atom is replaced by an aryl radical as defined above, such as, for example, benzyl, 2-phenylethyl and the like.

20       The term “alkylaryl,” alone or in combination, refers to an aryl radical as defined above in which one H atom is replaced by an alkyl radical as defined above, such as, for example, tolyl, xylyl and the like.

The terms haloalkyl, haloalkenyl, haloalkynyl and haloalkoxy include alkyl, alkenyl, alkynyl and alkoxy structures, as described above, that are substituted with one or more fluorines, chlorines, bromines or iodines, or with combinations thereof.

The terms cycloalkyl, aryl, arylalkyl, heteroaryl, alkyl, alkynyl, alkenyl, haloalkyl and heteroalkyl include optionally substituted cycloalkyl, aryl, arylalkyl, heteroaryl, alkyl, alkynyl, alkenyl, haloalkyl and heteroalkyl groups.

5 The term "carbocycle" includes optionally substituted, saturated or unsaturated, three- to eight-membered cyclic structures in which all of the skeletal atoms are carbon.

The term "heterocycle" includes optionally substituted, saturated or unsaturated, three- to eight-membered cyclic structures in which one or more skeletal atoms is oxygen, nitrogen, sulfur, phosphorus or combinations thereof. Illustrative examples include pyridine, pyran, thiophan, pyrrole, furan, thiophen, pentatomic and hexatomic lactam  
10 rings, and the like.

The term "membered ring" can embrace any cyclic structure, including carbocycles and heterocycles as described above. The term "membered" is meant to denote the number of skeletal atoms that constitute the ring. Thus, for example, pyridine, pyran, and thiophan are 6 membered rings and pyrrole, furan, and thiophen are 5  
15 membered rings.

The term "acyl" includes alkyl, aryl, heteroaryl, arylalkyl or heteroarylalkyl substituents attached to a compound via a carbonyl functionality (*e.g.*, -CO-alkyl, -CO-aryl, -CO-arylalkyl or -CO-heteroarylalkyl, etc.).

"Optionally substituted" groups may be substituted or unsubstituted. The  
20 substituents of an "optionally substituted" group may include, without limitation, one or more substituents independently selected from the following groups or designated subsets thereof: alkyl, alkenyl, alkynyl, heteroalkyl, haloalkyl, haloalkenyl, haloalkynyl, cycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, alkoxy, aryloxy, haloalkoxy, amino, alkylamino, dialkylamino, alkylthio, arylthio, heteroarylthio, oxo, carboxyesters, carboxamido, acyloxy, halogens, CN, NO<sub>2</sub>, NH<sub>2</sub>, N<sub>3</sub>, NHCH<sub>3</sub>, N(CH<sub>3</sub>)<sub>2</sub>, SH, SCH<sub>3</sub>, OH,  
25 OCH<sub>3</sub>, OCF<sub>3</sub>, CH<sub>3</sub>, CF<sub>3</sub>, C(O)CH<sub>3</sub>, CO<sub>2</sub>CH<sub>3</sub>, CO<sub>2</sub>H, C(O)NH<sub>2</sub>, , pyridinyl, thiophene, furanyl, indole, indazol, esters, amides, phosphonates, phosphates, phosphoramides, sulfonates, sulfates, sulphonamides, carbamates, ureas, thioureas, thioamides, thioalkyls. An optionally substituted group may be unsubstituted (*e.g.*, -CH<sub>2</sub>CH<sub>3</sub>), fully substituted

(*e.g.*, -CF<sub>2</sub>CF<sub>3</sub>), monosubstituted (*e.g.*, -CH<sub>2</sub>CH<sub>2</sub>F) or substituted at a level anywhere in-between fully substituted and monosubstituted (*e.g.*, -CH<sub>2</sub>CF<sub>3</sub>).

The term "halogen" includes F, Cl, Br and I.

5 The term sulfide refers to a sulfur atom covalently linked to two atoms; the formal oxidation state of said sulfur is (II). The term "thioether" may used interchangeably with the term "sulfide".

The term "sulfoxide" refers to a sulfur atom covalently linked to three atoms, at least one of which is an oxygen atom; the formal oxidation state of said sulfur atom is (IV).

10 The term "sulfone" refers to a sulfur atom covalently linked to four atoms, at least two of which are oxygen atoms; the formal oxidation state of said sulfur atom is (VI).

15 Some of the compounds of the present invention may contain one or more chiral centers and therefore may exist in enantiomeric and diastereomeric forms. The scope of the present invention is intended to cover all isomers per se, as well as mixtures of cis and trans isomers, mixtures of diastereomers and racemic mixtures of enantiomers (optical isomers) as well. Further, it is possible using well known techniques to separate the various forms, and some embodiments of the invention may feature purified or enriched species of a given enantiomer or diastereomer.

20 A "pharmacological composition" refers to a mixture of one or more of the compounds described herein, or pharmaceutically acceptable salts thereof, with other chemical components, such as pharmaceutically acceptable carriers and/or excipients. The purpose of a pharmacological composition is to facilitate administration of a compound to an organism.

25 The phrase "pharmaceutically acceptable carrier" as used herein means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the subject agent from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible

with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations. A physiologically acceptable carrier should not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered compound.

An "excipient" refers to an inert substance added to a pharmacological composition to further facilitate administration of a compound. Examples of excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

A "pharmaceutically effective amount" means an amount which is capable of providing a therapeutic and/or prophylactic effect. The specific dose of compound administered according to this invention to obtain therapeutic and/or prophylactic effect will, of course, be determined by the particular circumstances surrounding the case, including, for example, the specific compound administered, the route of administration, the condition being treated, and the individual being treated. A typical daily dose (administered in single or divided doses) will contain a dosage level of from about 0.01 mg/kg to about 50-100 mg/kg of body weight of an active compound of the invention. Preferred daily doses generally will be from about 0.05 mg/kg to about 20 mg/kg and ideally from about 0.1 mg/kg to about 10 mg/kg. Factors such as clearance rate and half-life and maximum tolerated dose (MTD) have yet to be determined but one of ordinary skill in the art can determine these using standard procedures.

In some method embodiments, the preferred therapeutic effect is the inhibition, to some extent, of the growth of cells characteristic of a proliferative disorder, e.g., breast cancer. A therapeutic effect will also normally, but need not, relieve to some extent one or more of the symptoms other than cell growth or size of cell mass. A therapeutic effect  
5 may include, for example, one or more of 1) a reduction in the number of cells; 2) a reduction in cell size; 3) inhibition (i.e., slowing to some extent, preferably stopping) of cell infiltration into peripheral organs, e.g., in the instance of cancer metastasis; 3) inhibition (i.e., slowing to some extent, preferably stopping) of tumor metastasis; 4) inhibition, to some extent, of cell growth; and/or 5) relieving to some extent one or more  
10 of the symptoms associated with the disorder.


In some method embodiments of the invention, the "IC<sub>50</sub>" value of a compound of the invention can be greater for normal cells than for cells exhibiting a proliferative disorder, e.g., breast cancer cells. The value depends on the assay used.

By a "standard" is meant a positive or negative control. A negative control in the  
15 context of HER-2 expression levels is, e.g., a sample possessing an amount of HER-2 protein that correlates with a normal cell. A negative control may also include a sample that contains no HER-2 protein. By contrast, a positive control does contain HER-2 protein, preferably of an amount that correlates with overexpression as found in proliferative disorders, e.g., breast cancers.. The controls may be from cell or tissue  
20 samples, or else contain purified ligand (or absent ligand), immobilized or otherwise. In some embodiments, one or more of the controls may be in the form of a diagnostic "dipstick."

By "selectively targeting" is meant affecting one type of cell to a greater extent than another, e.g., in the case of cells with high as opposed to relatively low or normal  
25 Her-2 levels.

### **Synthesis of the Compounds of the Invention**

The following synthesis scheme 1 is applicable to various of the compounds, compositions, methods, and formulations of the invention:



Scheme 1

Synthesis of compounds of formula 1 (when X = C) in scheme 1 may include some or all of the following general steps. The 8-substituted purine analogs of formula 5 or 2 can be prepared from 4,5-diaminopyrimidines and the carboxylates or their derivatives, such as amides, esters, nitriles, orthoesters, imidates etc (see, e.g., Townsend Chemistry of Nucleosides and Nucleotides, Vol. 1; Plenum Press, New York and London, page 148-158; Tetrahedron Lett. 36, 4249, 1995). Substituted 4,5-diaminopyrimidines can be obtained commercially or from substituted 2-chloro-3-amino pyrimidine or 2-chloro-3-nitropyrimidines as known in the art. See, e.g., Tetrahedron, 40, 1433 (1984); J. Am. Chem. Soc., 118, 135 (1975); Synthesis 135 (1975); J. Med. Chem. 39, 4099 (1996).

Compounds of formula 5 can be converted to compounds of formula 2 by simple alkylation with alkylhalides, alkyltosylates, mesolates or triflates in polar solvents like THF, DMF or DMSO using bases like NaH, Cs<sub>2</sub>CO<sub>3</sub> or K<sub>2</sub>CO<sub>3</sub>, or by the well-known Mitsunobu alkylation method.

Compounds of formula 2 can be further modified to give compounds of formula 1 or the intermediates to prepare compounds of formula 1, e.g., substitution of 6-chloropurine by ammonia or alkylamines. C-2 substitution of purines, e.g., halogenation with F, Cl or Br can be introduced via 2-aminopurines as described by Eaton et al., J. Org. Chem. 34(3), 747-8 (1969) or by nucleophilic substitution as described, e.g., in J. Med. Chem. 36, 2938 (1993) and Heterocycles, 30, 435, (1990). These C-2 substitutions also

can be introduced via metalation as described, e.g., in J. Org. Chem. 62(20), 6833 (1997), followed by addition of desired electrophile. General purine substitution can be accomplished as described in J. Med. Chem. 42, 2064 (1999).


Alternatively, intermediates of formula 2 can be prepared from  
5 chloroaminopyrimidines such as formula 6 by the following two steps: (1) treatment of the compounds of formula 6 with corresponding amine (Y-NH<sub>2</sub>), e.g., butylamine, in presence of base such as triethyl amine or N,N-diisopropyl amine in polar solvents such as n-BuOH to give the substituted diamine compounds of formula 4; (2) treatment of the  
10 compounds of formula 4 using the same methods as described earlier going from formula 7 to formula 5. Similar methods as described earlier can be used to introduce the C-2 substitution (point at which Z or G moiety attaches).

Compounds of formula 1 where A is other than NH<sub>2</sub>, e.g., halogen, methoxy, alkyl, or trifluoro alkyl, can be prepared starting with the corresponding substituent in place (if it can withstand the transformations), or, for halogen or substituted amines, these can be  
15 prepared from the 6-amine.

The compounds of formula 1 can also be prepared from formula 3, where L is halogen, using Negishi-type couplings (e.g., as described in J. Org. Chem. 2001, 66, 7522; J. Org. Chem. 1991, 56, 1445).

Compounds of formula 1 wherein X is a heteroatom such as S, O or N can be  
20 prepared by the following scheme 2. In general, these compounds are linked via their C-8 to one of the heteroatoms X = S, O, or N and can be prepared from the corresponding 8-halo (e.g., bromo, iodo or chloro) compounds such as formula 10 using nucleophiles such as sulfides, alkyl or arylthiols, amines, azides, and alcohols.





Scheme 2

With reference to scheme 2, substituted adenines or purines of formula 8 can be treated with halogenating agents such as bromine or iodine, followed by alkylation at N-9 to give compounds of formula 10, wherein M is halogen such as bromine or iodine (Dang et.al. PCT, WO 98/39344). Compounds of formula 16 can be prepared from trihalopyrimidines such as those of formula 12 by nitration to give compounds of formula 13. Subsequent displacement of the halogen with amine (YNH<sub>2</sub>) and reduction of the nitrogroup gives the diamines of formula 15. Alternatively, reduction of the nitrogroup may precede halogen displacement. Diamines of formula 15 can be readily cyclized to the imidazole ring of the compounds of formula 16, wherein L is H, SH, OH or NH<sub>2</sub> (Org. Syn. Collective Vol. 2, 65; Org. Syn. Collective Vol. 4, 569). The compounds of formula 1 can also be synthesized from the compounds of formula 16, wherein L is SH, OH, or NH<sub>2</sub>, by reacting with aromatic halides, boronic acids, triflates, or their equivalents in presence of a catalyst such as palladium or copper (Buchwald, S. L. et. al. J. Am. Chem. Soc., **1998**, *120*, 213-214; Buchwald, S. L. et. al. Acc. Chem. Res. **1998**, *31*, 805; Buchwald, S. L. et. al. Org. Lett., **2002**, *4*, 3517-3520).

Alternately, compounds of formulae 1 and 11 (wherein X= S or O) can be synthesized by coupling of the diazonium salts of the compounds of formulae 10 or 16 (wherein M or L is N<sub>2</sub>.BF<sub>4</sub>, N<sub>2</sub>.HCl, N<sub>2</sub>.H<sub>2</sub>SO<sub>4</sub> etc.) with HXE or HXQ (wherein X=S or O) in the presence of base such as t-BuOK, NaH, etc. in solvents such as DMF, MeOH, etc.

Z-groups of formula 1 can be introduced by modifying existing 2-substituents such as G. For example, 2-halopurines of formula 1 can be prepared from 2-aminopurines (G = NH<sub>2</sub>) via chemistry well described in the literature. Other substitutions such as S-alkyl or aryl, O-alkyl can be made from nucleophilic substitution reactions; metal-catalysed reactions, etc. (see, e.g., Aerschot et. al., J .Med. Chem. 36:2938 (1993); Buchwald, S. L. et. al., Heterocycles, 30: 435 (1990).

The E component (aromatic or heteroaromatic or alkyl) of the compounds of formula 11 can be further modified as needed using well known procedures including, e.g., nucleophilic additions, electrophilic additions, halogenations, etc. to give Q (see, e.g., Advanced Organic Chemistry, March. J. Wiley Interscience).

Compounds of formula 1, wherein X is S(O) or S(O)<sub>2</sub> can be prepared by the oxidation of the compounds of formula 1, wherein X=S, using reagents such as MCPBA, H<sub>2</sub>O<sub>2</sub>, NaIO<sub>4</sub>, Oxone, etc. in solvents such as CHCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub> etc. Also, these sulfone compounds can be made by coupling of sulfonyl salts such as Li, Na, K (ArS(O)<sub>2</sub>Li) and compounds of formulae 10 or 16 (wherein M or L is halogen such as Br or I) in polar solvents such as DMF. (Chem. Abstr. 1952, 4549). With controlled reduction of these sulfones, one can make compounds of formula 1 where X is S(O) and S(O)<sub>2</sub>.

### Pharmaceutical Compositions, Dosaging and Modes of Administration

Those of ordinary skill in the art are familiar with formulation and administration techniques that can be employed with the compounds and methods of the invention, e.g., as discussed in Goodman and Gilman's, *The Pharmacological Basis of Therapeutics*, current edition; Pergamon Press; and Remington's *Pharmaceutical Sciences* (current edition.) Mack Publishing Co., Easton, Pa.

The compounds utilized in the methods of the instant invention may be administered either alone or in combination with pharmaceutically acceptable carriers, excipients or diluents, in a pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

For example, the therapeutic or pharmaceutical compositions of the invention can be administered locally to the area in need of treatment. This may be achieved by, for example, but not limited to, local infusion during surgery, topical application, e.g., cream, ointment, injection, catheter, or implant, said implant made, e.g., out of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. The administration can also be by direct injection at the site (or former site) of a tumor or neoplastic or pre-neoplastic tissue.

Still further, the compounds or compositions of the invention can be delivered in a vesicle, e.g., a liposome (see, for example, Langer, **1990**, *Science*, 249:1527-1533; Treat et al., **1989**, *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Bernstein and Fidler (eds.), Liss, N.Y., pp. 353-365).

The compounds and pharmaceutical compositions used in the methods of the present invention can also be delivered in a controlled release system. In one embodiment, a pump may be used (see, Sefton, **1987**, *CRC Crit. Ref. Biomed. Eng.* 14:201; Buchwald et al., **1980**, *Surgery*, 88:507; Saudek et al., **1989**, *N. Engl. J. Med.*, 321:574). Additionally, a controlled release system can be placed in proximity of the therapeutic target. (see, Goodson, **1984**, *Medical Applications of Controlled Release*, Vol. 2, pp. 115-138).

The pharmaceutical compositions used in the methods of the instant invention can also contain the active ingredient in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions, and such compositions may contain one or more agents

selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets.

5 These excipients may be, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, such as microcrystalline cellulose, sodium crosscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may  
10 be un-coated or coated by known techniques to mask the taste of the drug or delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water soluble taste masking material such as hydroxypropylmethyl-cellulose or hydroxypropylcellulose, or a time delay material such as ethyl cellulose, or cellulose acetate butyrate may be employed as appropriate.

15 Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

20 Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide,  
25 for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethylene-oxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with  
30 partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more

preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

5 Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated  
10 hydroxyanisol or alpha-tocopherol.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already  
15 mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

The compounds and pharmaceutical compositions used in the methods of the instant invention may also be in the form of an oil-in-water emulsions. The oily phase  
20 may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphatides, for example soy bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example  
25 polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening agents, flavoring agents, preservatives and antioxidants.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, flavoring and coloring agents and antioxidant.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous solutions. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

5 The sterile injectable preparation may also be a sterile injectable oil-in-water microemulsion where the active ingredient is dissolved in the oily phase. For example, the active ingredient may be first dissolved in a mixture of soybean oil and lecithin. The oil solution then introduced into a water and glycerol mixture and processed to form a microemulsion.

10 The injectable solutions or microemulsions may be introduced into a patient's blood-stream by local bolus injection. Alternatively, it may be advantageous to administer the solution or microemulsion in such a way as to maintain a constant circulating concentration of the instant compound. In order to maintain such a constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

15 The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension for intramuscular and subcutaneous administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in  
20 a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

25 The Compounds of the present invention used in the methods of the present invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the inhibitors with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials  
30 include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of

polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing an compound or composition of the invention can be used. As used herein,  
5 topical application can include mouth washes and gargles.

The compounds used in the methods of the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery  
10 system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

The methods, compounds and compositions of the instant invention may also be used in conjunction with other well known therapeutic agents that are selected for their particular usefulness against the condition that is being treated. For example, the instant  
15 compounds may be useful in combination with known anti-cancer and cytotoxic agents. Further, the instant methods and compounds may also be useful in combination with other inhibitors of parts of the signaling pathway that links cell surface growth factor receptors to nuclear signals initiating cellular proliferation.

The methods of the present invention may also be useful with other agents that  
20 inhibit angiogenesis and thereby inhibit the growth and invasiveness of tumor cells, including, but not limited to VEGF receptor inhibitors, including ribozymes and antisense targeted to VEGF receptors, angiostatin and endostatin.

Examples of antineoplastic agents that can be used in combination with the compounds and methods of the present invention include, in general, and as appropriate,  
25 alkylating agents, anti-metabolites; epidophyllotoxin; an antineoplastic enzyme; a topoisomerase inhibitor; procarbazine; mitoxantrone; platinum coordination complexes; biological response modifiers and growth inhibitors; hormonal/anti-hormonal therapeutic agents and haematopoietic growth factors. Exemplary classes of antineoplastics include the anthracycline family of drugs, the vinca drugs, the mitomycins, the bleomycins, the  
30 cytotoxic nucleosides, the epothilones, discodermolide, the pteridine family of drugs,

diylenes and the podophyllotoxins. Particularly useful members of those classes include, for example, carminomycin, daunorubicin, aminopterin, methotrexate, methopterin, dichloromethotrexate, mitomycin C, porfiromycin, 5-fluorouracil, 6-mercaptopurine, gemcitabine, cytosine arabinoside, podophyllotoxin or podo-phyllotoxin derivatives such as etoposide, etoposide phosphate or teniposide, melphalan, vinblastine, vincristine, leurosine, vindesine, leurosine, paclitaxel and the like. Other useful antineoplastic agents include estramustine, carboplatin, cyclophosphamide, bleomycin, gemcitabine, ifosamide, melphalan, hexamethyl melamine, thiotepa, cytarabin, idatrexate, trimetrexate, dacarbazine, L-asparaginase, camptothecin, CPT-11, topotecan, ara-C, bicalutamide, flutamide, leuprolide, pyridobenzoindole derivatives, interferons and interleukins.

When a compound or composition of the invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for cancer, for example, breast cancer. Administration typically occurs in an amount of between about 0.01 mg/kg of body weight to about 100 mg/kg of body weight per day (administered in single or divided doses), more preferably at least about 0.1 mg/kg of body weight per day. A particular therapeutic dosage can include, e.g., from about 0.01 mg to about 1000 mg of compound, and preferably includes, e.g., from about 1 mg to about 1000 mg. The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 0.1 mg to 1000 mg, preferably from about 1 mg to 300 mg, more preferably 10 mg to 200 mg, according to the particular application. The amount administered will vary depending on the particular IC<sub>50</sub> value of the compound used and the judgment of the attending clinician taking into consideration factors such as health, weight, and age. In combinational applications in which the compound is not the sole active ingredient, it may be possible to administer lesser amounts of compound and still have therapeutic or prophylactic effect.

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose.



The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound.

5 Thereafter, the dosage is increased by small amounts until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

The amount and frequency of administration of the compounds and compositions of the present invention used in the methods of the present invention, and if applicable  
10 other chemotherapeutic agents and/or radiation therapy, will be regulated according to the judgment of the attending clinician (physician) considering such factors as age, condition and size of the patient as well as severity of the disease being treated.

The chemotherapeutic agent and/or radiation therapy can be administered according to therapeutic protocols well known in the art. It will be apparent to those  
15 skilled in the art that the administration of the chemotherapeutic agent and/or radiation therapy can be varied depending on the disease being treated and the known effects of the chemotherapeutic agent and/or radiation therapy on that disease. Also, in accordance with the knowledge of the skilled clinician, the therapeutic protocols (e.g., dosage amounts and times of administration) can be varied in view of the observed effects of the administered  
20 therapeutic agents (i.e., antineoplastic agent or radiation) on the patient, and in view of the observed responses of the disease to the administered therapeutic agents.

Also, in general, the compounds of the invention need not be administered in the same pharmaceutical composition as a chemotherapeutic agent, and may, because of different physical and chemical characteristics, be administered by a different route. For  
25 example, the compounds/compositions may be administered orally to generate and maintain good blood levels thereof, while the chemotherapeutic agent may be administered intravenously. The determination of the mode of administration and the advisability of administration, where possible, in the same pharmaceutical composition, is well within the knowledge of the skilled clinician. The initial administration can be made  
30 according to established protocols known in the art, and then, based upon the observed

effects, the dosage, modes of administration and times of administration can be modified by the skilled clinician.

The particular choice of compound (and where appropriate, chemotherapeutic agent and/or radiation) will depend upon the diagnosis of the attending physicians and their judgment of the condition of the patient and the appropriate treatment protocol.

The compounds/compositions of the invention (and where appropriate chemotherapeutic agent and/or radiation) may be administered concurrently (e.g., simultaneously, essentially simultaneously or within the same treatment protocol) or sequentially, depending upon the nature of the proliferative disease, the condition of the patient, and the actual choice of chemotherapeutic agent and/or radiation to be administered in conjunction (i.e., within a single treatment protocol) with the compound/composition.

In combinational applications and uses, if the compound/composition and the chemotherapeutic agent and/or radiation are not administered simultaneously or essentially simultaneously, then the initial order of administration of the compound/composition, and the chemotherapeutic agent and/or radiation, may not be important. Thus, the compounds/compositions of the invention may be administered first followed by the administration of the chemotherapeutic agent and/or radiation; or the chemotherapeutic agent and/or radiation may be administered first followed by the administration of the compounds/compositions of the invention. This alternate administration may be repeated during a single treatment protocol. The determination of the order of administration, and the number of repetitions of administration of each therapeutic agent during a treatment protocol, is well within the knowledge of the skilled physician after evaluation of the disease being treated and the condition of the patient. For example, the chemotherapeutic agent and/or radiation may be administered first, especially if it is a cytotoxic agent, and then the treatment continued with the administration of the compounds/compositions of the invention followed, where determined advantageous, by the administration of the chemotherapeutic agent and/or radiation, and so on until the treatment protocol is complete.

Thus, in accordance with experience and knowledge, the practicing physician can modify each protocol for the administration of a compound/composition for treatment according to the individual patient's needs, as the treatment proceeds.

5 The attending clinician, in judging whether treatment is effective at the dosage administered, will consider the general well-being of the patient as well as more definite signs such as relief of disease-related symptoms, inhibition of tumor growth, actual shrinkage of the tumor, or inhibition of metastasis. Size of the tumor can be measured by standard methods such as radiological studies, e.g., CAT or MRI scan, and successive measurements can be used to judge whether or not growth of the tumor has been retarded  
10 or even reversed. Relief of disease-related symptoms such as pain, and improvement in overall condition can also be used to help judge effectiveness of treatment.

#### **Assays to Determine HSP90 Binding and Downstream Effect**

A variety of in vitro and in vivo assays are available to test the effect of the compounds of the invention on HSP90. HSP90 competitive binding assays and functional  
15 assays can be performed as known in the art substituting in the compounds of the invention. Chiosis et al., Chemistry & Biology 8:289-299 (2001), describe some of the known ways in which this can be done. For example, competition binding assays using, e.g., geldanamycin or 17-AAG as a competitive binding inhibitor of HSP90 can be used to determine relative HSP90 affinity of the compounds of the invention by immobilizing the  
20 compound of interest or other competitive inhibitor on a gel or solid matrix, preincubating HSP90 with the other inhibitor, passing the preincubated mix over the gel or matrix, and then measuring the amount of HSP90 that sticks or does not stick to the gel or matrix.

Downstream effects can also be evaluated based on the known effect of HSP90 inhibition on function and stability of various steroid receptors and signaling proteins  
25 including, e.g., Raf1 and Her2. Compounds of the present invention induce dose-dependent degradation of these molecules, which can be measured using standard techniques. Inhibition of HSP90 also results in up-regulation of HSP90 and related chaperone proteins that can similarly be measured. Antiproliferative activity on various cancer cell lines can also be measured, as can morphological and functional differentiation  
30 related to HSP90 inhibition. For example, the

Many different types of methods are known in the art for determining protein concentrations and measuring or predicting the level of proteins within cells and in fluid samples. Indirect techniques include nucleic acid hybridization and amplification using, e.g., polymerase chain reaction (PCR). These techniques are known to the person of skill and are discussed, e.g., in Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., Ausubel, et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, 1994, and, as specifically applied to the quantification, detection, and relative activity of Her-2/neu in patient samples, e.g., in U.S. Patents 4,699,877, 4,918,162, 4,968,603, and 5,846,749. A brief discussion of two generic techniques that can be used follows.

The determination of whether cells overexpress or contain elevated levels of HER-2 can be determined using well known antibody techniques such as immunoblotting, radioimmunoassays, western blotting, immunoprecipitation, enzyme-linked immunosorbant assays (ELISA), and derivative techniques that make use of antibodies directed against HER-2. As an example, HER-2 expression in breast cancer cells can be determined with the use of an immunohistochemical assay, such as the Dako Hercep™ test (Dako Corp., Carpinteria, CA). The Hercep™ test is an antibody staining assay designed to detect HER-2 overexpression in tumor tissue specimens. This particular assay grades HER-2 expression into four levels: 0, 1, 2, and 3, with level 3 representing the highest level of HER-2 expression. Accurate quantitation can be enhanced by employing an Automated Cellular Imaging System (ACIS) as described, e.g., by Press, M, et al, (2000), *Modern Pathology* 13:225A.

Antibodies, polyclonal or monoclonal, can be purchased from a variety of commercial suppliers, or may be manufactured using well-known methods, e.g., as described in Harlow et al., *Antibodies: A Laboratory Manual*, 2nd Ed; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1988).

HER-2 overexpression can also be determined at the nucleic acid level since there is a reported high correlation between overexpression of the HER-2 protein and amplification of the gene that codes for it. One way to test this is by using RT-PCR. The genomic and cDNA sequences for HER-2 are known. Specific DNA primers can be

generated using standard, well-known techniques, and can then be used to amplify template already present in the cell. An example of this is described in Kurokawa, H et al, Cancer Res. 60: 5887-5894 (2000). PCR can be standardized such that quantitative differences are observed as between normal and abnormal cells, e.g., cancerous and noncancerous cells. Well known methods employing, e.g., densitometry, can be used to quantitate and/or compare nucleic acid levels amplified using PCR.

Similarly, fluorescent in situ hybridization (FISH) assays and other assays can be used, e.g., Northern and/or Southern blotting. These rely on nucleic acid hybridization between the HER-2 gene or mRNA and a corresponding nucleic acid probe that can be designed in the same or a similar way as for PCR primers, above. See, e.g., Mitchell MS, and Press MF., 1999, Semin. Oncol., Suppl. 12:108-16. For FISH, this nucleic acid probe can be conjugated to a fluorescent molecule, e.g., fluorescein and/or rhodamine, that preferably does not interfere with hybridization, and which fluorescence can later be measured following hybridization. See, e.g., Kurokawa, H et al, Cancer Res. 60: 5887-5894 (2000) (describing a specific nucleic acid probe having sequence 5'-FAM-NucleicAcid-TAMRA-p-3' sequence). ACIS-based approaches as described above can be employed to make the assay more quantitative (de la Torre-Bueno, J, et al , 2000, Modern Pathology 13:221A).

Immuno and nucleic acid detection can also be directed against proteins other than HSP90 and Her-2, which proteins are nevertheless affected in response to HSP90 inhibition.

The following examples are offered by way of illustration only and are not intended to be limiting of the full scope and spirit of the invention.

### EXAMPLES

The chemical reagents used below are all available commercially, e.g., from Aldrich Chemical Co., Milwaukee, WI, USA, and/or their facile preparation known to one of ordinary skill in the art, or otherwise described or referenced herein.

**Carbon linker compounds:****Example 1:** Synthesis of 8-(2,5-dimethoxybenzyl)-N9-butyladenine

**Step 1.** A solution of 5-amino-4,6-dichloropyrimidine (1 mmol) in n-BuOH was treated with Et<sub>3</sub>N (1.2 mmol) and n-Butylamine (1.0 mmol) at 80 °C. After 16 h, solvent was removed under reduced pressure. The residue was dissolved in EtOAc, the organic layer washed with water and then dried (MgSO<sub>4</sub>). Filtration and removal of solvent gave 6-chloro-5-amino-4-butyl pyrimidine as a brown solid. R<sub>f</sub> = 0.5 in 1:1 EtOAc:hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.07 (s, 1H), 4.88 (br s, 1H), 3.49 (m, 2H), 3.35 (br s, 2H), 1.6 (m, 2H), 1.44 (m, 2H), 0.95 (t, 3H).

**Step 2:** To a solution of 2,5-dimethoxyphenylacetic acid (1 mmol) and Et<sub>3</sub>N (1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was added p-toluenesulfonyl chloride (1 mmol) at rt. After 1 h, the mixture was treated with a solution of the product of step 1, 6-chloro-5-amino-4-butyl pyrimidine (1 mmol in CH<sub>2</sub>Cl<sub>2</sub>), followed by addition of Et<sub>3</sub>N (2 mmol). The resultant mixture was refluxed for 20 h. Solvent was removed and the residue dissolved into EtOAc, the organic layer washed with water and dried. The crude compound was taken into acetone, and precipitated product filtered out and washed with a small amount of acetone to give N-(4-butylamino-6-chloro-pyrimidin-5-yl)-2-(2,5-dimethoxyphenyl) acetamide. R<sub>f</sub> = 0.45 in 1:1 EtOAc:hexane. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.37 (s, 1H), 8.17 (s, 1H), 7.11 (t, 1H), 6.9 (d, 1H), 6.88 (d, 1H), 6.78 (dd, 1H), 3.73 (s, 3H), 3.69 (s, 3H), 3.63 (s, 3H), 3.35 (m, 2H), 1.48 (m, 2H), 1.29 (m, 2H), 0.88 (t, 3H).

**Step 3:** A mixture of N-(4-butylamino-6-chloro-pyrimidin-5-yl)-2-(2,5-dimethoxyphenyl) acetamide (1 mmol) and p-TSA (0.5 mmol) in toluene was refluxed for 72 h. Solvent was removed, diluted with EtOAc and washed with water, bicarbonate and dried. Purification on a silica gel column (200-400 mesh, Fisher Scientific, Tustin, CA, USA) gave 6-chloro-8-(2,5-dimethoxybenzyl)-N9-butyl purine. R<sub>f</sub> = 0.65 in 1:1 EtOAc:hexane. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.7 (s, 1H), 6.96 (d, 1H), 6.84 (m, 1H), 6.8 (dd, 1H), 4.28 (s, 2H), 4.23 (t, 2H), 3.69 (s, 3H), 3.67 (s, 3H), 1.62 (m, 2H), 1.25 (m, 2H), 0.88 (t, 3H).

**Step 4:** To a solution of 6-chloro-8-(2,5-dimethoxybenzyl)-N9-butyl purine (1 mmol) in dioxane was added 28 % NH<sub>4</sub>OH (50 mmol) and the mixture was then heated at 100 °C in a seal tube for 48 h. Solvent was removed by azeotrope distillation with toluene. Purification on a silica gel column (see above) gave pure 8-(2,5-dimethoxybenzyl)-9-butyl adenine, **1.1**. R<sub>f</sub> = 0.35 in 5 % MeOH in EtOAc. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.08 (s, 1H), 7.04 (br s, 2H), 6.94 (d, 1H), 6.80 (dd, 1H), 6.66 (d, 1H), 4.14 (s, 2H), 4.04 (t, 2H), 3.72 (s, 3H), 3.63 (s, 3H), 1.52 (m, 2H), 1.22 (m, 2H), 0.82 (t, 3H).

Alternatively, 8-(2,5-dimethoxybenzyl)-9-butyl adenine can also be prepared from N-(4-butylamino-6-chloro-pyrimidin-5-yl)-2-(2,5-dimethoxyphenyl) acetamide according to the following procedure: A solution of N-(4-butylamino-6-chloro-pyrimidin-5-yl)-2-(2,5-dimethoxyphenyl) acetamide (1 mmol) is taken into 7M NH<sub>3</sub> in MeOH (70 mmol) and the mixture heated at 120 °C in a steel bomb for 72 h. Solvent is removed by azeotrope distillation with toluene. Purification on the silica gel column gave pure 8-(2,5-dimethoxybenzyl)-9-butyl adenine.

**Example 2: Synthesis of 8-(2,5-dimethoxybenzyl)-N9-pentynyl-2-fluoro adenine**

**Step 1: 2-(2,5-Dimethoxy-phenyl)-N-(2,5,6-triamino-pyrimidin-4-yl)-acetamide, HCl**

A solution of 2,4,5,6-tetraaminopyrimidine (52.8 g, 378 mmol) in NMP (750 ml) was treated at 70 °C with 2,5-dimethoxyphenyl acetyl chloride (90 g, 419 mmol). After cooling to r.t., the precipitate was collected by filtration and washed with EtOAc to give the title compounds as a pale yellow powder (127 g, 95%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.12 (s, 1H), 7.80–7.40 (m, 3H), 6.22 (s, 2H), 6.04 (s, 4H), 4.41 (s, 3H), 4.29 (s, 3H), 4.25 (s, 2H); MS 319 (M+1).

**Step 2: 8-(2,5-Dimethoxy-benzyl)-9H-purine-2,6-diamine**

Sodium metal (2.3 g, 100 mmol) was dissolved in n-BuOH (50 ml) at 70 °C. To this was added the acetamide of step 1, above (5.0 g, 14.1 mmol), and the mixture was heated to reflux for 1.5 h. Neutralization with 6N HCl to pH 8–9, extraction with EtOAc, drying, and evaporation gave the title compound as a pale yellow powder (3.2 g, 76%). R<sub>f</sub> = 0.45 in 1:3 MeOH:EtOAc. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 12.3–11.7 (br. s, 1H), 6.92 (d, J =

10.0 Hz, 1H), 6.82 (dd, J = 10.0 & 3.0 Hz, 1H), 6.73 (s, 1H), 6.70–6.50 (br. s, 2H), 5.85–5.70 (br. s, 2H), 3.95 (s, 2H), 3.74 (s, 3H), 3.67 (s, 3H); MS 301 (M+1).

**Step 3: 8-(2,5-Dimethoxy-benzyl)-9-pent-4-ynyl-9H-purine-2,6-diamine:**

A mixture of the purine 8-(2,5-Dimethoxy-benzyl)-9H-purine-2,6-diamine (19.0 g, 63 mmol), 5-chloro-pent-1-yne (12.3 ml, 116 mmol), and Cs<sub>2</sub>CO<sub>3</sub> (37.8 g, 116 mmol) in DMF (180 g) was heated to 50 °C for 16h. Filtration and washing (2 x 200 ml H<sub>2</sub>O) afforded some desired product (5.8 g, 25%). The mother liquor was concentrated, diluted with EtOAc, and heated to reflux for 1h to yield additional product (6.0 g, 26%). After cooling to room temperature, addition of 1 volume hexane to the EtOAc mother liquor gave additional product (2.6 g, 11%). Final work-up (CH<sub>2</sub>Cl<sub>2</sub>:MeOH 4:1 – water) yielded additional product (5.3 g, containing 1 equivalent penty-4-yn-1-ol, 18%). R<sub>f</sub> = 0.65 in 1:10 MeOH:EtOAc. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 6.92 (d, J = 8.9 Hz, 1H), 6.98 (dd, J = 8.9 & 3.0 Hz, 1H), 6.59 (s, J = 2.9 Hz, 1H), 6.58–6.53 (br. s, 2H), 5.72–5.68 (br. s, 2H), 4.02 (s, 2H), 3.92 (t, J = 7.4 Hz, 2H), 3.73 (s, 3H), 3.62 (s, 3H), 2.84 (t, J = 2.5 Hz, 1H), 2.13 (td, J = 7.0 & 1.7 Hz, 2H), 1.74 (quint., J = 7.3 Hz, 2H); MS 367 (M+1).

**Step 4: 8-(2,5-Dimethoxy-benzyl)-2-fluoro-9-pent-4-ynyl-9H-purin-6-ylamine**

A solution of the above purine-2,6-diamine (11.8 g, 32.2 mmol) in 48% aq. HBF<sub>4</sub> (250 ml) was treated at –10 °C with iso-amyl nitrite (5.20 ml, 38.8 mmol), and warmed to r.t over 2.5 h. The reaction mixture was diluted with MeOH (400 ml) and CH<sub>2</sub>Cl<sub>2</sub> (1500 ml), and carefully neutralized with a solution of K<sub>2</sub>CO<sub>3</sub> (125 g) in water (500 ml). Caution: vigorous gas evolution. The aqueous layer was re-extracted with MeOH:CH<sub>2</sub>Cl<sub>2</sub> (500 ml, 1:5). Concentration of the organic phase and two flash chromatography purifications (CH<sub>2</sub>Cl<sub>2</sub>:EtOAc:hexane:MeOH:Et<sub>3</sub>N 1500:750:750:50:10 → 1500:750:750:150:10) yielded 8-(2,5-Dimethoxy-benzyl)-2-fluoro-9-pent-4-ynyl-9H-purin-6-ylamine (4.5 g, 38%), 2.1 as a colorless powder. R<sub>f</sub> = 0.45 in 1:1 EtOAc:hexane. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 6.82 (d, J = 8.9 Hz, 1H), 6.75 (dd, J = 8.9 & 3.0 Hz, 1H), 6.68 (d, J = 2.9 Hz, 1H), 6.25–6.10 (br. s, 2H), 4.20 (s, 2H), 4.13 (t, J = 7.4 Hz, 2H), 3.79 (s, 3H), 3.70 (s, 3H), 2.16 (td, J = 7.0 & 2.6 Hz, 2H), 1.97 (t, J = 2.6 Hz, 1H), 1.95 (quint., J = 7.3 Hz, 2H); MS 370 (M+1).



The following compounds, 2.2-2.4, were prepared using essentially the same procedures described for Example 2, except that in step 3 the electrophiles 1-bromo-4-methyl-pent-3-ene, 1-chloro-pent-4-ene, and 1,5-bromopentane were used in place of 5-chloro-pent-1-yne:

5    **2.2    8-(2,5-Dimethoxy-benzyl)-2-fluoro-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine** isolated as solid, retention time = 7.70.

**2.3    8-(2,5-Dimethoxy-benzyl)-2-fluoro-9-pent-4-enyl-9H-purin-6-ylamine** isolated as solid, retention time = 7.61.

10    **2.4    8-(2,5-Dimethoxy-benzyl)-2-fluoro-9-(5-bromo-pentyl)-9H-purin-6-ylamine** isolated as solid, retention time = 7.86.

Similarly, 2-Cl compound was prepared analogously to the method described in Step 4 using HCl and CuCl in place of HBF<sub>4</sub>.

**2.5    8-(2,5-Dimethoxy-benzyl)-2-chloro-9-pent-4-ynyl-9H-purin-6-ylamine;**

15    Rt=8.02 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.83 (d, J = 8.9 Hz, 1H), 6.77 (dd, J = 8.9 & 3.0 Hz, 1H), 6.68 (d, J = 3.0 Hz, 1H), 6.18–6.00 (s, 2H), 4.20 (s, 2H), 4.18 (t, J = 7.4 Hz, 2H), 3.78 (s, 3H), 4.93 (s, 3H), 2.20 (td, J = 7.0 & 2.4 Hz, 2H), 2.63 (t, 2.4 Hz, 1H), 1.97 (quint., J = 7.3 Hz, 2H).

HPLC method: Agilent Zorbax 300 SB C18, 4.6 X 150 mm, 5 μm; Column Temperature: Ambient; Flow Rate: 1.0 ml/min, Gradient: 10% acetonitrile (0.05% TFA) in water (0.1% TFA) to 100% acetonitrile (0.05% TFA) in 10 minutes, hold at 100 % for 1 minutes);  
20    Retention times are measured in minutes.

**Example 3:** The above procedures can similarly be applied to produce compounds wherein Z is H by starting with 4,5,6, triaminopyrimidine sulfate and using electrophiles as shown in Table 3:

Table 3

Ex. #	Electrophile	Final Compound/Structure/Name/HPLC RT (min.)/CF#
3.1	1-bromo-4-chlorobutane	9-(4-Chloro-butyl)-8-(2,5-dimethoxy-benzyl)-9H-purin-6-ylamine isolated as solid; rt = 6.34.
3.2	1-chloro-pent-4-yne	8-(2,5-Dimethoxy-benzyl)-9-pent-4-ynyl-9H-purin-6-ylamine isolated as solid rt= 5.88 min.
3.3	1-chloro-3-[1,3]dioxolan-2-yl]propane	8-(2,5-Dimethoxy-benzyl)-9-(2-[1,3]dioxolan-2-yl-ethyl)-9H-purin-6-ylamine isolated as solid, rt = 5.36.
3.4	1-bromo-4-methyl-pent-3-ene	8-(2,5-Dimethoxy-benzyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine isolated as solid., rt = 6.60.
3.5	1,5-dibromopentane	9-(5-Bromo-pentyl)-8-(2,5-dimethoxy-benzyl)-9H-purin-6-ylamine isolated as solid, rt = 6.94.
3.6	1,5-dibromo-3-methylpentane	9-(5-Bromo-3-methyl-pentyl)-8-(2,5-dimethoxy-benzyl)-9H-purin-6-ylamine isolated as solid, rt = 7.32.
3.7	1-bromo-5-Chloropentane	9-(5-Chloro-pentyl)-8-(2,5-dimethoxy-benzyl)-9H-purin-6-ylamine isolated as solid, rt = 6.34.
3.8	1-bromo-4-chlorobutane alkylation followed by treatment with ethylamine gave 4-ethylaminobutyl	8-(2,5-Dimethoxy-benzyl)-9-(4-ethylamino-butyl)-9H-purin-6-ylamine isolated as solid, rt = 3.9.
3.9	1-bromohexan-6-ol	6-[6-Amino-8-(2,5-dimethoxy-benzyl)-purin-9-yl]-hexan-1-ol isolated as solid.
3.10	2-(dimethyl-bicyclo[3.1.1]hept-2-en-2-yl)-3-bromopropane	8-(2,5-Dimethoxy-benzyl)-9-[2-(dimethyl-bicyclo[3.1.1]hept-2-en-2-yl)-ethyl]-9H-purin-6-ylamine isolated as solid.
3.11	1-bromo-5acetyloxypentane	Acetic acid 5-[6-amino-8-(2,5-dimethoxy-benzyl)-purin-9-yl]-pentyl ester isolated as solid; rt =6.06.
3.12	1-bromo-3,3,3-trifluoro-	8-(2,5-Dimethoxy-benzyl)-9-(3,3,3-trifluoro-propyl)-9H-purin-

	propane	6-ylamine isolated as solid;
3.13	1-chloro-pent-4-yne	8-(2,5-Dimethoxy-benzyl)-9-pent-4-ynyl-9H-purin-6-ylamine isolated as solid; rt = 5.88.

#### Example 4: Halogenation of benzene ring

4.1 To a solution of 9-butyl-8-(3-methoxy-benzyl)-9H-purin-6-ylamine (1.24 g, 4 mmol) in AcOH (6 ml) was added N-iodo-succinamide (NIS) (1.8 g, 8 mmol). After 3h at r.t., additional NIS (1.8 g, 8 mmol) was added, and the mixture was stirred for another 24 h. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (500 ml), and carefully neutralized with a solution of sat. aq. K<sub>2</sub>CO<sub>3</sub> (2 x 100 ml), then washed with 0.1 N Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (3 x 100 ml), brine (3 x 100 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated, and purified by flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>: MeOH=100:5) to give the 9-Butyl-8-(2-iodo-5-methoxy-benzyl)-9H-purin-6-ylamine (**4.1**), as a colorless powder (0.53 g, 30%); rt = 7.7 min.; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d) δ 8.36 (s, 1H), 7.77 (d, J = 7.9 Hz, 1H), 6.68 (s, 1H), 6.61 (d, J = 7.9 Hz, 1H), 5.62 (s, 2H), 4.33 (s, 2H), 4.06 (t, J = 7.7 Hz, 2H), 3.72 (s, 3H), 1.67 (quint., J = 7.7 Hz, 2H), 1.36 (sext., J = 7.5 Hz, 2H), 0.92 (t, J = 7.4 Hz, 3H).

Bromo and chloro derivatives were made using the same procedure, substituting NBS and NCS for NIS as appropriate. The following compounds were also synthesized according to essentially the same procedure, using as appropriate NIS, NCS or NBS:

4.2 **9-Butyl-8-(5-iodo-2-methoxy-benzyl)-9H-purin-6-ylamine** was made from 9-Butyl-8-(2-methoxy-benzyl)-9H-purin-6-ylamine as starting material in 48 % yield; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.32 (s, 1H), 7.55 (dd, J = 8.7, 2.2 Hz, 1H), 7.37 (d, J = 2.2 Hz, 1H), 6.68 (d, J = 8.7 Hz, 1H), 6.05–5.85 (br. s, 2H), 4.17 (s, 2H), 4.07 (t, J = 7.6 Hz, 2H), 3.82 (s, 3H), 1.62 (quint., J = 7.5 Hz, 2H), 1.30 (sext., J = 7.5 Hz, 2H), 0.89 (t, J = 7.4 Hz, 3H).

4.3 **9-Butyl-8-(5-ethyl-2-methoxy-benzyl)-9H-purin-6-ylamine**; Rt = 7.59; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d) δ 8.35 (s, 1H), 7.34 (d, J = 8.8 Hz, 1H), 6.79 (dd, J = 8.7, 2.8 Hz, 1H), 6.69 (d, J = 2.7 Hz, 1H), 5.64 (s, 2H), 4.36 (s, 2H), 4.07 (t, J = 7.7 Hz, 2H), 3.73 (s, 3H), 1.64 (quint., J = 7.7 Hz, 2H), 1.32 (sext., J = 7.5 Hz, 2H), 0.90 (t, J = 7.4 Hz, 3H).

4.4 **8-(2-Bromo-5-methoxy-benzyl)-9-butyl-9H-purin-6-ylamine;** Rt = 7.66; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d) δ 8.36 (s, 1H), 7.52 (d, J = 8.7 Hz, 1H), 6.74 (dd, J = 8.7, 3.0 Hz, 1H), 6.89 (d, J = 3.0 Hz, 1H), 5.64 (s, 2H), 4.36 (s, 2H), 4.07 (t, J = 7.7 Hz, 2H), 3.72 (s, 3H), 1.64 (quint., J = 7.6 Hz, 2H), 1.34 (sext., J = 7.5 Hz, 2H), 0.90 (t, J = 7.4 Hz, 3H).

5 9-Butyl-8-(2-methoxy-benzyl)-9H-purin-6-ylamine and 9-butyl-8-(3-methoxy-benzyl)-9H-purin-6-ylamine were prepared from 4,5,6-triaminopyrimidine sulfate and, respectively 2-methoxyphenyl acetyl chloride or 3-methoxyphenyl acetic acid, by procedures analogous to the one described above. 2-Fluoro purine analogs were also prepared from 2,4,5,6-tetraaminopyrimidine, by procedures analogous to those described  
10 above. See Example 2, step 4.

For the following compounds, in which the N9 substituent (Y) is sensitive to halogenation, addition of the N9 substituent (Y) was done as a final step:

4.5 **8-(2-Bromo-5-methoxy-benzyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine;** Rt = 8.22; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d) δ 8.37 (s, 1H), 7.51 (d, J = 8.7 Hz, 1H), 6.73 (dd, J = 8.7  
15 Hz, 3.0 Hz, 1H), 6.65 (d, J = 3.0 Hz, 1H), 5.53 (s, 2H), 5.12 (t, J = 7.1 Hz, 2H), 4.35 (s, 2H), 4.07 (t, J = 7.1 Hz, 2H), 3.72 (s, 3H), 2.43 (quart., J = 7.1 Hz, 2H), 1.65 (s, 3H), 1.40 (s, 3H).

4.6 **8-(2-Bromo-5-methoxy-benzyl)-9-pent-4-ynyl-9H-purin-6-ylamine;** Rt = 8.17; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d) δ 8.35 (s, 1H), 7.52 (d, J = 8.8 Hz, 1H), 6.74 (dd, J = 8.8 Hz, 2.9 Hz,  
20 1H), 6.66 (d, J = 2.9 Hz, 1H), 5.61 (s, 2H), 4.39 (s, 2H), 4.21 (t, J = 7.4 Hz, 2H), 3.73 (s, 3H), 2.24 (td, J = 6.8 Hz, 2.5 Hz, 2H), 2.03 (t, J = 2.5 Hz, 1H), 1.99 (quint., J = 7.2 Hz, 2H).

4.7 **8-(2-Iodo-5-methoxy-benzyl)-9-pent-4-ynyl-9H-purin-6-ylamine;** Rt = 7.35; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d) δ 8.36 (s, 1H), 7.77 (d, J = 8.5 Hz, 1H), 6.64-6.60 (m, 2H), 5.56 (s,  
25 2H), 4.35 (s, 2H), 4.20 (t, J = 7.4 Hz, 2H), 3.73 (s, 3H), 2.26 (td, J = 6.9 Hz, 2.7 Hz, 2H), 2.03 (t, J = 2.7 Hz, 1H), 2.02 (quint., J = 7.0 Hz, 2H).

4.8 **8-(2-Iodo-5-methoxy-benzyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine;** Rt = 8.17; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d) δ 8.58 (s, 1H), 8.33 (d, J = 8.6 Hz, 1H), 6.60 (d, J = 2.9 Hz,

1H), 6.57 (dd, J = 8.6, 2.9 Hz, 1H), 6.15 (s, 2H), 5.12 (t, J = 7.4 Hz, 2H), 4.29 (s, 2H), 4.04 (t, J = 7.3 Hz, 2H), 3.67 (s, 3H), 2.42 (quart., J = 7.2 Hz, 2H), 1.65 (s, 3H), 1.39 (s, 3H).

**4.9 2-Fluoro-8-(2-iodo-5-methoxy-benzyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine;** Rt = 10.04; <sup>1</sup>H NMR (CDCl<sub>3</sub>-d) δ 7.76 (d, J = 8.6 Hz, 1H), 6.65 (d, J = 2.5 Hz, 1H), 6.60 (dd, J = 8.6, 2.5 Hz, 1H), 6.14 (s, 2H), 5.13 (t, J = 6.9 Hz, 1H), 4.26 (s, 2H), 4.01 (t, J = 7.0 Hz, 2H), 3.72 (s, 3H), 2.43 (quint., J = 7.0 Hz, 2H), 1.68 (s, 3H), 1.42 (s, 3H).

**4.10 2-Fluoro-8-(2-iodo-5-methoxy-benzyl)-9-pent-4-ynyl-9H-purin-6-ylamine;** Rt = 8.75; <sup>1</sup>H NMR (CDCl<sub>3</sub>-d) δ 7.77 (d, J = 8.7 Hz, 1H), 6.67 (d, J = 2.7 Hz, 1H), 6.62 (dd, J = 8.7, 2.7 Hz, 1H), 5.99 (s, 2H), 4.32 (s, 2H), 4.16 (t, J = 7.2 Hz, 2H), 3.74 (s, 3H), 2.26 (td, J = 6.7, 2.6 Hz, 2H), 2.02 (t, J = 2.4 Hz, 1H), 1.99 (quint., J = 6.9 Hz, 2H); MP: 172-177°C.

**Example 5: General procedure for palladium-mediated couplings:**

A mixture of 9-Butyl-8-(5-iodo-2-methoxy-benzyl)-9H-purin-6-ylamine (50 mg, 0.1 mmol) and Pd(PPh<sub>3</sub>)<sub>4</sub> (12 mg, 0.01 mmol) was treated under N<sub>2</sub> at r.t. with a 1M solution of the organometallic coupling partner (0.5 ml, 0.5 mmol). Reactions were performed typically in THF at r.t. for 10 min with organomagnesium compounds in THF at r.t. for 16 h with organozinc compounds, or in DMF at 80 °C for 3h with organostannanes. After work-up, the product was purified by chromatography on preparative plates (1000 uM, SiO<sub>2</sub>), eluting with CH<sub>2</sub>Cl<sub>2</sub>:EtOAc:hexane: MeOH: Et<sub>3</sub>N 1500:750:750:50:10.

The following compounds were prepared using the corresponding commercially available organozinc compound; the skilled artisan will recognize that equivalent organostannane, and organoboron, and organomagnesium coupling partners may be used in place of organozinc compounds. A general review of appropriate methodologies may be found in "Palladium Reagents in Organic Synthesis" Richard F. Heck, Academic Press, 1990.

**5.1 9-Butyl-8-(5-ethyl-2-methoxy-benzyl)-9H-purin-6-ylamine;** Rt = 8.23; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.30 (s, 1H), 7.07 (dd, J = 8.4 & 2.0 Hz, 1H), 6.91 (d, J = 2.0 Hz, 1H), 6.83 (d, J = 8.4 Hz, 1H), 5.65-5.55 (s, 2H), 4.23 (s, 2H), 4.04 (t, J = 7.6 Hz, 2H),

3.83 (s, 3H), 2.51 (q, J = 7.6 Hz, 2H) 1.65–1.55 (m, 2H), 1.30–1.25 (m, 2H), 1.41 (t, J = 7.6 Hz, 3H), 0.86 (t, J = 7.3 Hz, 3H).

**5.2 9-Butyl-8-(5-butyl-2-methoxy-benzyl)-9H-purin-6-ylamine;** Rt = 9.24; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.33 (s, 1H), 7.05 (dd, J = 8.4 & 1.9 Hz, 1H), 6.88 (d, J = 1.8 Hz, 1H), 6.82 (d, J = 8.3 Hz, 1H), 5.58–5.48 (s, 2H), 4.23 (s, 2H), 4.04 (t, J = 7.6 Hz, 2H), 3.83 (s, 3H), 2.47 (q, J = 7.6 Hz, 2H), 1.57 (quint., J = 7.5 Hz, 2H), 1.48 (quint., J = 7.6 Hz, 2H), 1.32–1.22 (m, 4H), 0.87 (t, J = 7.3 Hz, 3H), 0.86 (t, J = 7.3 Hz, 3H).

**5.3 9-Butyl-8-(2-methoxy-5-vinyl-benzyl)-9H-purin-6-ylamine;** Rt = 7.91; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.31 (s, 1H), 7.31 (dd, J = 8.5 & 2.3 Hz, 1H), 7.16 (d, J = 2.2 Hz, 1H), 6.87 (d, J = 8.5 Hz, 1H), 6.59 (dd, J = 17.6 & 10.9 Hz, 1H), 5.82–5.72 (s, 2H), 5.53 (dd, J = 17.6 & 0.7 Hz, 1H), 5.09 (dd, J = 10.9 & 0.7 Hz, 1H), 4.22 (s, 2H), 4.06 (t, J = 7.6 Hz, 2H), 3.85 (s, 3H), 1.62 (quint., J = 7.7 Hz, 2H), 1.30 (sext., J = 7.4 Hz, 2H), 0.87 (t, J = 7.4 Hz, 3H).

**Example 6: General procedure for the nitration of benzene ring and**

**derivatizations:**

A solution of purine in H<sub>2</sub>SO<sub>4</sub> or in H<sub>2</sub>SO<sub>4</sub>:AcOH 1:4 was treated at 0 °C with 1 equiv HNO<sub>3</sub>. The mixture was diluted with EtOAc, neutralized with NaHCO<sub>3</sub> and purified by chromatography on SiO<sub>2</sub> preparative plates (1000 uM) with CH<sub>2</sub>Cl<sub>2</sub>: EtOAc:hexane: MeOH: Et<sub>3</sub>N 1500:750:750:50:10.

Nitro derivatives (20 mg) are reduced with 10%Pd/C (Aldrich) (20 mg) under H<sub>2</sub> atmosphere in THF at r.t. over 16h. The resulting aniline can be further monoalkylated (Acetylchloride, CH<sub>2</sub>Cl<sub>2</sub>) or reductively alkylated (RCHO, NaBH(OAc)<sub>3</sub>, 1,2-dichloroethane, r.t.) .

The following compounds were prepared by this method:

**6.1 8-(2,5-Dimethoxy-4-nitro-benzyl)-2-fluoro-9-pent-4-ynyl-9H-purin-6-ylamine** (CF310), Rt = 8.05; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.94 (s, 1H), 6.85 (s, 1H), 6.37–6.27 (s, 2H), 4.06 (s, 2H), 4.01 (t, J = 7.3 Hz, 2H), 3.69 (s, 3H), 3.66 (s, 3H), 2.13 (td, J = 7.0 & 2.6 Hz, 2H), 1.87 (t, J = 2.6 Hz, 1H), 1.82 (quint., J = 7.3 Hz, 2H).

**6.2 9-Butyl-8-(3,5-dimethoxy-2-nitro-benzyl)-9H-purin-6-ylamine; sulfuric acid salt;** Rt = 7.33;  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  8.27 (s, 1H), 8.15–7.90 (br. s, 2H), 6.78 (d, J = 2.4 Hz, 1H), 6.55 (d, J = 2.4 Hz, 1H), 4.32 (s, 2H), 4.12 (t, J = 7.3 Hz, 2H), 3.88 (s, 3H), 3.81 (s, 3H), 1.58 (quint., J = 7.5 Hz, 2H), 1.21 (sext., J = 7.5 Hz, 2H), 0.84 (t, J = 7.4 Hz, 3H).

**6.3 8-(4-Amino-3,5-dimethoxy-benzyl)-9-butyl-9H-purin-6-ylamine;** Rt = 8.05;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.31 (s, 1H), 7.31 (dd, J = 8.5 & 2.3 Hz, 1H), 7.16 (d, J = 2.2 Hz, 1H), 6.87 (d, J = 8.5 Hz, 1H), 6.59 (dd, J = 17.6 & 10.9 Hz, 1H), 5.82–5.72 (s, 2H), 5.53 (dd, J = 17.6 & 0.7 Hz, 1H), 5.09 (dd, J = 10.9 & 0.7 Hz, 1H), 4.22 (s, 2H), 4.06 (t, J = 7.6 Hz, 2H), 3.85 (s, 3H), 1.62 (quint., J = 7.7 Hz, 2H), 1.30 (sext., J = 7.4 Hz, 2H), 0.87 (t, J = 7.4 Hz, 3H).

**6.4 8-(4-Amino-2,5-dimethoxy-benzyl)-9-butyl-9H-purin-6-ylamine;** Rt = 6.95;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.33 (s, 1H), 6.57 (s, 1H), 6.33 (s, 1H), 6.37–6.27 (s, 2H), 4.20 (s, 2H), 4.01 (t, J = 7.3 Hz, 2H), 3.74 (s, 3H), 3.68 (s, 3H), 1.59 (quint., J = 7.5 Hz, 2H), 1.32 (sext., J = 7.5 Hz, 2H), 0.86 (t, J = 7.4 Hz, 3H).

**6.5 8-(2-Amino-3,5-dimethoxy-benzyl)-9-butyl-9H-purin-6-ylamine;**  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.28 (s, 1H), 6.40 (d, J = 2.5 Hz, 1H), 6.30 (d, J = 2.5 Hz, 1H), 5.85–5.75 (s, 2H), 4.14 (s, 2H), 4.13 (t, J = 7.6 Hz, 2H), 3.80 (s, 3H), 3.73 (s, 3H), 1.62 (quint., J = 7.5 Hz, 2H), 1.48 (sext., J = 7.5 Hz, 2H), 0.91 (t, J = 7.4 Hz, 3H).

**6.6 2-(6-Amino-9-butyl-9H-purin-8-ylmethyl)-4-methoxy-benzaldehyde-O-methyl-oxime;** Rt = 7.69;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.88 (s, 1H), 8.31 (s, 1H), 7.72 (d, J = 7.9 Hz, 1H), 6.80 (d, J = 8.0 Hz, 1H), 6.74 (s, 1H), 5.80–5.76 (s, 2H), 4.24 (s, 2H), 4.00 (t, J = 7.7 Hz, 2H), 3.94 (s, 3H), 3.76 (s, 3H), 1.58 (quint., J = 7.7 Hz, 2H), 1.28 (sext., J = 7.5 Hz, 2H), 0.86 (t, J = 7.3 Hz, 3H).

#### **Example 7: Formylation of benzene ring and derivatization**

A solution of 9-butyl-8-(3-methoxy-benzyl)-9H-purin-6-ylamine (100 mg, 0.32 mmol), 1,1-dichlorodimethyl ether (40 mg, 0.35 mmol) and  $\text{TiCl}_4$  (133 mg, 0.70 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 ml) was prepared at  $0^\circ\text{C}$  and stirred at r.t. overnight. Dilution with  $\text{CH}_2\text{Cl}_2$ ,

washing ( $\text{Na}_2\text{SO}_4$ ,  $\text{NH}_4\text{Cl}$ ), drying, and preparative thin layer chromatography gave the title aldehyde as a yellow glass (47 mg, 43%).

Standard procedures gave the corresponding alcohol ( $\text{NaBH}_4$ , MeOH, r.t.), tosyl hydrazone ( $\text{TsNHNH}_2$ , EtOH, reflux), oximes ( $\text{RONH}_2 \cdot \text{HCl}$ , DMF, 60 °C), amines (5  $(\text{R}_1\text{R}_2\text{NH}$ ,  $\text{NaBH}(\text{OAc})_3$ ,  $\text{Cl}-(\text{CH}_2)_2\text{-Cl}$  r.t.), homoallylic alcohol ( $\text{AlSiMe}_3$ ,  $\text{TiCl}_4$ ),  $\text{CH}_2\text{Cl}_2$ , -78 °C), or alkenes.

7.1 **2-(6-Amino-9-butyl-9H-purin-8-ylmethyl)-4-methoxy-benzaldehyde;** Rt = 6.52;  $^1\text{H}$ NMR ( $\text{CDCl}_3\text{-d}$ )  $\delta$  10.39 (s, 1H), 8.32 (s, 1H), 7.76 (d, J = 7.8 Hz, 1H), 6.87 (m, 2H), 6.22 (s, 2H), 4.28 (s, 2H), 4.03 (t, J = 7.6 Hz, 2H), 3.85 (s, 3H), 1.61 (quint., J = 7.3 10 Hz, 2H), 1.29 (sext., J = 7.4 Hz, 2H), 0.86 (t, J = 7.2 Hz, 3H).

#### Example 8: Negishi couplings

A mixture of 3,4-dichlorobenzyl bromide (0.47 g, 1.96 mmol) and Rieke Zinc (3.0 ml, 5g/100 ml THF, 2.35 mmol) was stirred overnight at r.t. in a flame-dried Schlenk tube and decanted to provide a 0.65M stock solution of 3,4-dichlorobenzyl zinc bromide. A 15 solution of 8-bromo-9-butyl-9H-purin-6-ylamine (42.7 mg, 0.158 mol),  $\text{Pd}(\text{dppf})\text{Cl}_2$  (16.8 mg, 0.020 mmol), and 3,4-dichlorobenzyl zinc bromide (0.61 ml, 0.65M in THF) was stirred in a flame-dried Schlenk tube at 66°C overnight, quenched with sat. aq.  $\text{NH}_4\text{Cl}$  and sat. aq. EDTA., extracted into EtOAc, dried and concentrated. Preparative TLC purification (EtOAc/ $\text{CH}_2\text{Cl}_2$ /MeOH 14:14:2) provided the title compound as a colorless 20 oil (approx. 15 mg, 20%).

8.1 **9-Butyl-8-(3,4-dichloro-benzyl)-9H-purin-6-ylamine,** compound isolated as solid, Rt = 7.98.

#### S- Linker compounds

##### Example 9: 9-Butyl-8-(2-iodo, 5-methoxy-phenylsulfanyl)-9H-purin-6-ylamine

25 **Step 1:** Adenine (47 g, 0.35 mole) was suspended in 200 ml of  $\text{CHCl}_3$  before adding bromine (180 ml, 3.5 mole) in one portion. The suspension was left stirring at room temperature for 72 hours in a closed system that was vented by a 20G needle. The reaction was worked up by adding shaved ice into the suspension before slowly



neutralizing with aqueous ammonia to pH 8-9, followed by precipitation of the desired product with acetic acid. The crude product was dried under reduced pressure for 2 days to give 8-Bromoadenine as a light brown powder (45 g, 60% yield).  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  8.12 (s, 1H), 7.22 (s, 2H). Rf(75% EtOAc/Hex) = 0.4.

- 5 **Step 2:** 8-Bromopurine (2.2g, 10mmole) was dissolved in 50 ml of DMF before adding 1-bromo-butane (2.2 ml, 20 mmol) and cesium carbonate (6.7g, 20 mmol) into the solution. The reaction mixture was left stirring at room temperature for 16 hours before quenching with water and extracting with EtOAc. The organic layer was washed with water and dried with  $\text{MgSO}_4$  before removing solvent under reduced pressure. A white powder (0.9g, 33%) of 8-Bromo-9-butyl-9H-purin-6-ylamine was isolated using silica gel column chromatography (50% EtOAc/Hexanes).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.32, (s, 1H), 5.81 (s, 2H), 4.20 (t, 2H), 1.82 (m, 2H), 1.40 (m, 2H), 0.96 (t, 3H). Rf(75% EtOAc/Hex) = 0.6.

- 15 **Step 3:** To a mixture of sodium hydride (96 mg, 4 mmol) in DMF (4ml) was added 3-methoxy-benzenethiol (1.12 g, 8 mmol). After 30 min, a solution of 8-bromo-9-butyl-9H-purin-6-ylamine (0.54 g, 2 mmol) in DMF (6 ml) was added and stirred for 12 h at 70 °C. The reaction mixture was quenched by addition of MeOH (4ml), diluted with EtOAc (400 ml), washed with  $\text{Na}_2\text{CO}_3$  (3  $\times$  100 ml), brine (3  $\times$  100 ml), dried ( $\text{Na}_2\text{SO}_4$ ), evaporated, purified with flash chromatography ( $\text{CH}_2\text{Cl}_2$ :MeOH = 100:5) to give the title sulfide as a colorless powder (0.59 g, 89%).

- 20 HPLC method used for these compounds: Agilent Zorbax 300 SB C18, 4.6 X 150 mm, 5  $\mu\text{m}$ ; Column Temperature: Ambient; Flow Rate: 1.0 ml/min, Gradient: 5% acetonitrile (0.05% TFA) in water (0.1% TFA) to 100% acetonitrile (0.05% TFA) in 15 minutes, hold at 100 % for 2 minutes).

- 25 The following compounds were prepared using the corresponding thiol in place of the 3-methoxybenzene thiol used in step 3:

**9.1 3-(6-Amino-9-butyl-9H-purin-8-ylsulfanyl)-phenol;** Rt= 6.75 min  $^1\text{H}$ NMR (DMSO- $d_6$ ):  $\delta$  9.69 (s, 1H), 8.17 (s, 1H), 7.45 (s, 2H), 7.17 (t, J=7.9Hz, 1H), 6.76(d, J=7.4Hz, 1H), 6.68 (d, J=8.2Hz, 1H), 6.62 (s, 1H), 4.11 (t, J=7.0Hz, 2H), 1.57 (quint., J=7.3Hz, 2H), 1.19 (sext., J=6.8Hz, 2H), 0.81 (t, J=7.4Hz, 3H).

**9.2 9-Butyl-8-(3-methoxy-phenylsulfanyl)-9H-purin-6-ylamine;** Rt= 8.6 min; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.80 (t, J = 7.4 Hz, 3H, CH<sub>3</sub>), 1.20 (m, 2H, CH<sub>2</sub>), 1.61 (m, 2H, CH<sub>2</sub>), 3.60 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 4.13 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H).

**9.3 9-Butyl-8-(2,5-dimethoxy-phenylsulfanyl)-9H-purin-6-ylamine;** Rt= 7.62 min; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d): δ 8.30 (s, 1H), 7.18 (t, J=8.2Hz, 1H), 6.90 (m, 2H), 6.77 (m, 3H), 4.17 (t, J=7.6Hz, 2H), 3.70 (s, 3H), 1.67 (quint., J=7.5Hz, 2H), 1.28 (sext., J=7.5Hz, 2H), 0.86 (t, J=7.4Hz, 3H).

**Step 4:** To a solution of 9-butyl-8-(3-methoxy-phenylsulfanyl)-9H-purin-6-ylamine (0.26g, 0.73mmol) in AcOH ( 4ml) was added NIS (0.53g, 2.19mmol) in portions. The mixture was stirred for 24 h at r.t. The reaction mixture was diluted with EtOAc (200 ml), and carefully neutralized with a solution of K<sub>2</sub>CO<sub>3</sub> (2 × 50 ml), then washed with Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (3 × 50 ml), brine (3 × 50 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated, purified by preparative TLC chromatography (CH<sub>2</sub>Cl<sub>2</sub>: MeOH = 100:5) to give the 2-iodo isomer (60 mg), and the 4-iodo isomer (65 mg).

**9.4 9-Butyl-8-(2-iodo-5-methoxy-phenylsulfanyl)-9H-purin-6-ylamine;** Rt= 8.45 min; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d): δ 8.38 (s, 1H), 7.73 (d, J=8.7Hz, 1H), 6.71 (d, J=2.7Hz, 1H), 6.58 (dd, J=8.7, 2.7Hz, 1H), 5.91 (s, 2H), 4.22 (t, J=7.4Hz, 2H), 3.68 (s, 3H), 1.75 (quint., J=7.7Hz, 2H), 1.34 (sext., J=7.5Hz, 2H), 0.93 (t, J=7.4Hz, 3H).

**9.5 9-Butyl-8-(4-iodo-3-methoxy-phenylsulfanyl)-9H-purin-6-ylamine;** Rt= 8.63 min; <sup>1</sup>HNMR (CDCl<sub>3</sub>-d): δ 8.38 (s, 1H), 7.72 (d, J=8.1Hz, 1H), 6.92 (d, J=1.8Hz, 1H), 6.58 (dd, J=8.1, 1.8Hz, 1H), 5.82 (s, 2H), 4.22 (t, J=7.4Hz, 2H), 3.85 (s, 3H), 1.75 (quint., J=7.7Hz, 2H), 1.37 (sext., J=7.5Hz, 2H), 0.93 (t, J=7.4Hz, 3H).

For compounds in which the N9 substituent (Y) is sensitive to halogenation conditions, these may be prepared using iodide already present in the benzenethiol moiety:

To a suspension of sodium hydride (96 mg, 4 mmol) in DMF (3 ml) was added 2-iodo-5-methoxy-benzenethiol (1.06 g, 4 mmol; J Org. Chem, 2001, 66(13), 4525-4542). After 30 min, a solution of 8-bromo-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine (296

mg, 1 mmol) in DMF (3 ml) was added, and the mixture was stirred for 12 h at 70 °C. The reaction was quenched by addition of MeOH (2 ml), diluted with EtOAc (200 ml), washed with Na<sub>2</sub>CO<sub>3</sub> (3 × 50 ml), brine (3 × 50 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated, and purified by flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>: MeOH= 100:5) to give 8-(2-Iodo-5-methoxy-phenylsulfanyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine as a colorless powder (280 mg, 58%).

The following compounds were prepared by this method using, respectively, the electrophiles 1-bromo-4-methyl-pent-3-ene and 1-chloro-pent-4-yn:

**9.6 8-(2-Iodo-5-methoxy-phenylsulfanyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine;** Rt= 9.14 min; <sup>1</sup>H NMR (CDCl<sub>3</sub>-d): δ 8.39 (s, 1H), 7.72 (d, J=8.7Hz, 1H), 6.72 (d, J=2.7Hz, 1H), 6.58 (dd, J=8.7, 2.7Hz, 1H), 5.81 (s, 2H), 5.15 (t, J=7.3Hz, 1H), 4.25 (t, J=7.4Hz, 2H), 3.69 (s, 3H), 2.50 (quint., J=7.3Hz, 2H), 1.66 (s, 3H), 1.44 (s, 3H); MP: 167-167.5°C.

**9.7 8-(2-Iodo-5-methoxy-phenylsulfanyl)-9-pent-4-ynyl-9H-purin-6-ylamine;** Rt= 7.93 min; <sup>1</sup>H NMR (CDCl<sub>3</sub>-d): δ 8.38 (s, 1H), 7.75 (d, J=8.7Hz, 1H), 6.74 (d, J=2.7Hz, 1H), 6.60 (dd, J=8.7, 2.7Hz, 1H), 5.72 (s, 2H), 4.32 (t, J=7.3Hz, 2H), 3.70 (s, 3H), 2.28 (td, J=6.8, 2.6Hz, 2H), 2.06 (quint., J=7.3Hz, 2H), 2.00 (t, J=2.4Hz, 1H); MP: 168-169°C.

The following compounds were prepared using the corresponding thiol in place of the 3-methoxybenzene thiol and base t-BuOK in place of NaH used in step 3:

**9.8 8-(Benzothiazole-2-ylsulfanyl)-9-butyl-9H-purin-6-ylamine;** Rt=6.53 min; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 8.41 (s, 1H), 7.94 (d, 1H), 7.74 (d, 1H), 7.47 (t, 1H), 7.38 (t, 1H), 6.01 (s, 2H), 4.32 (t, 2H), 1.79 (m, 2H), 1.35 (m, 2H), 0.89 (t, 3H).

**9.9 9-Butyl-8-(5-chloro-benzothiazole-2-ylsulfanyl)-9H-purin-6-ylamine;**—Mass (M+1) = 391.8 et (M+3) = 393.8; <sup>1</sup>H NMR (CDCl<sub>3</sub>) 8.43 (s, 1H), 7.92 (s, 1H), 7.65 (d, 1H), 7.35 (d, 1H), 6.01 (s, 2H), 4.32 (t, 2H), 1.79 (m, 2H), 1.35 (m, 2H), 0.89 (t, 3H).

**9.10 9-Butyl-8-(5-methoxy-benzothiazole-2-ylsulfanyl)-9H-purin-6-ylamine;**—<sup>1</sup>H NMR (CDCl<sub>3</sub>) 8.42 (s, 1H), 7.60 (d, 1H), 7.43 (s, 1H), 7.02 (d, 1H), 5.82 (s, 2H), 4.33 (t, 2H), 3.99 (s, 3H), 1.80 (m, 2H), 1.35 (m, 2H), 0.89 (t, 3H).

**9.11 9-Butyl-8-(2,5-dichloro-phenylsulfanyl)-9H-purin-6-ylamine;**  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 8.37 (s, 1H), 7.35 (d, 1H), 7.20 (dd, 1H), 7.14 (d, 1H), 5.72 (s, 2H), 4.24 (t, 2H), 1.79 (m, 2H), 1.35 (m, 2H), 0.89 (t, 3H).

**9.12 9 Butyl-8-(2,4,5-trichloro-phenylsulfanyl)-9H-purin-6-ylamine;**  $R_t=7.8$  min;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 8.37 (s, 1H), 7.62 (s, 1H), 7.35 (s, 1H), 5.98 (s, 2H), 4.27 (t, 2H), 1.80 (m, 2H), 1.36 (m, 2H), 0.92 (t, 3H).

**Example 10:** 8-(2,5-dimethoxy-phenylsulfanyl)-2-fluoro-9(4-methyl-pent-3-enyl)-9H-purin-6-ylamine and 8-(2,5-dimethoxy-phenylsulfanyl)-2-amino-9(4-methyl-pent-3-enyl)-9H-purin-6-ylamine were prepared from 2,6-diaminopurine by procedures analogous to the one described above in Example 9. The final conversion of amino to fluoro was done by method similar to that reported in Example 2, step 4.

**10.1 8-(2,5-dimethoxy-phenylsulfanyl)-2-amino-9(4-methyl-pent-3-enyl)-9H-purin-6-ylamine;**  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  1.28 (s, 3H,  $\text{CH}_3$ ), 1.58 (s, 3H,  $\text{CH}_3$ ), 2.35 (m, 2H,  $\text{CH}_2$ ), 3.60 (s, 3H,  $\text{OCH}_3$ ), 3.76 (s, 3H,  $\text{OCH}_3$ ), 4.12 (t,  $J = 7.0$  Hz, 2H,  $\text{CH}_2$ ), 5.05 (t,  $J = 7$  Hz, 1H,  $\text{CH}=\text{)$ , 6.50 (s, 1H, Ar-H), 6.91 (d,  $J = 8.9$  Hz, 1H, Ar-H), 7.05 (d,  $J = 8.9$  Hz, 1H, Ar-H).

**10.2 8-(2,5-dimethoxy-phenylsulfanyl)-2-fluoro-9(4-methyl-pent-3-enyl)-9H-purin-6-ylamine;**  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  1.30 (s, 3H,  $\text{CH}_3$ ), 1.55 (s, 3H,  $\text{CH}_3$ ), 2.35 (m, 2H,  $\text{CH}_2$ ), 3.60 (s, 3H,  $\text{OCH}_3$ ), 3.76 (s, 3H,  $\text{OCH}_3$ ), 4.10 (t,  $J = 7.0$  Hz, 2H,  $\text{CH}_2$ ), 5.05 (t,  $J = 7$  Hz, 1H,  $\text{CH}=\text{)$ , 6.47 (s, 1H, Ar-H), 6.86 (d,  $J = 8.9$  Hz, 1H, Ar-H), 7.02 (d,  $J = 8.9$  Hz, 1H, Ar-H); MS ( $m/z$ ) 426 ( $\text{M}+\text{Na}$ ).

**Example 11:**

The compounds in this example were prepared analogously to the method described above in Example 9 using various electrophiles to generate a library of N9 substituted compounds. N9 alkylation was done as a final step after the bromine displacement of 8-bromopurine with 2,5-dimethoxy thiophenol.

**11.1 8-(2,5-dimethoxy-phenylsulfanyl)-9H-purin-6-ylamine;**  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  3.62 (s, 3H,  $\text{OCH}_3$ ), 3.76 (s, 3H,  $\text{OCH}_3$ ), 6.61 (s, 1H, Ar-H), 6.85 (d,  $J = 8.9$  Hz, 1H, Ar-

H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.24 (bs, 2H, NH<sub>2</sub>), 8.13 (s, 1H, purine-H) 13.33 (s, 1H, purine-NH); electrophile: No substitution on N9.

11.2 8-(2,5-dimethoxy-phenylsulfanyl)-9-pentyl-9H-purin-6-ylamine; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.80 (t, J = 7.4 Hz, 3H, CH<sub>3</sub>), 1.20 (m, 4H, 2CH<sub>2</sub>), 1.61 (m, 2H, CH<sub>2</sub>), 3.60 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 4.13 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H); electrophile: 1-bromopentyl.

11.3 8-(2,5-dimethoxy-phenylsulfanyl)-9-pent-4-ynyl-9H-purin-6-ylamine; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.89(m, 2H, CH<sub>2</sub>), 2.20(t, J=8.0Hz, 2H, CH<sub>2</sub>), 2.78(s, 1H, CH≡), 3.62 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 4.23 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H); electrophile: 1-chloro-pent-4-yne.

11.4 4-[6-Amino-8(2,5-dimethoxysulfanyl)-purin-9-yl]-butyronitrile; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.89(m, 2H, CH<sub>2</sub>), 2.20(t, J=8.0Hz, 2H, CH<sub>2</sub>), 3.62 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 4.23 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H); electrophile: 1-bromobutyronitrile.

11.5 8-(2,5-dimethoxy-phenylsulfanyl)-9(3,3,3-trifluoromethylpropyl)-9H-purin-6-ylamine; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.54(t, J=8.0Hz, 2H, CH<sub>2</sub>), 3.62 (s, 3H, OCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 4.46 (t, J = 8.0 Hz, 2H, CH<sub>2</sub>), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.30 (s, 1H, purine-H); electrophile: 1-bromo-3,3,3-trifluoro-propane.

11.6 8-(2,5-dimethoxy-phenylsulfanyl)-9(4-chlorobutyl)-9H-purin-6-ylamine; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.82(m, 2H, CH<sub>2</sub>), 1.98(m, 2H, CH<sub>2</sub>), 3.56(t, J=6.4Hz, 2H, CH<sub>2</sub>), 3.75 (s, 3H, OCH<sub>3</sub>), 3.78 (s, 3H, OCH<sub>3</sub>), 4.23 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H); electrophile: 1-bromo-4-chlorobutane.

11.7 8-(2,5-dimethoxy-phenylsulfanyl)-9(4-acetyloxybutyl)-9H-purin-6-ylamine; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.70(m, 2H, CH<sub>2</sub>), 1.90(m, 2H, CH<sub>2</sub>), 2.02(s, 3H, CH<sub>3</sub>), 3.75 (s, 3H,

OCH<sub>3</sub>), 3.78 (s, 3H, OCH<sub>3</sub>), 4.10 (t, J=6.4Hz, 2H, CH<sub>2</sub>), 4.30 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H); electrophile: 1-bromo-4-acetyloxybutane.

**11.8 8-(2,5-dimethoxy-phenylsulfanyl)-9(5-bromopentyl)-9H-purin-6-ylamine;** <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.46 (m, 2H, CH<sub>2</sub>), 1.85(m, 4H, 2 CH<sub>2</sub>), 3.36(t, J=6.7Hz, 2H, CH<sub>2</sub>), 3.72 (s, 3H, OCH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 4.30 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H); electrophile: 1,5-dibromopentane.

**11.9 8-(2,5-dimethoxy-phenylsulfanyl)-9(2-[1,3]dioxolan-2-yl-ethyl)-9H-purin-6-ylamine;** <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.26(m, 2H, CH<sub>2</sub>), 3.75 (s, 3H, OCH<sub>3</sub>), 3.77 (s, 3H, OCH<sub>3</sub>), 3.85(t, J=7.0Hz, 2H, CH<sub>2</sub>), 3.98(t, J=7.0Hz, 2H, CH<sub>2</sub>), 4.46 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 4.96(t, J=4.1Hz, 1H, CH), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H); electrophile: 2-(2-Chloro-ethyl)-[1,3]dioxolane.

**11.10 8-(2,5-dimethoxy-phenylsulfanyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine;** <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.28 (s, 3H, CH<sub>3</sub>), 1.54 (s, 3H, CH<sub>3</sub>), 2.35 (m, 2H, CH<sub>2</sub>), 3.60 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 4.15 (t, J = 7.0 Hz, 2H, CH<sub>2</sub>), 5.05 (t, J = 7 Hz, 1H, CH=), 6.46(s, 1H, Ar-H), 6.86 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.42(bs, 2H, NH<sub>2</sub>), 8.17 (s, 1H, purine-H); electrophile: 1-bromo-4-methyl-pent-3-ene; MP: 148-150°C.

**11.11 8-(2,5-dimethoxy-phenylsulfanyl)-9-(pent-4-enyl)-9H-purin-6-ylamine;** <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.89(m, 2H, CH<sub>2</sub>), 2.19(t, J=8.0Hz, 2H, CH<sub>2</sub>), 3.62 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 4.23 (t, J = 7.4 Hz, 2H, CH<sub>2</sub>), 5.05(m, 2H, CH<sub>2</sub>=), 5.82(m, 1H, CH=), 6.46(s, 1H, Ar-H), 6.85 (d, J = 8.9 Hz, 1H, Ar-H), 7.02 (d, J = 8.9 Hz, 1H, Ar-H), 7.41(bs, 2H, NH<sub>2</sub>), 8.15 (s, 1H, purine-H); electrophile: 1-chloro-pent-4-yne.

**11.12 8-(2,5-dimethoxy-phenylsulfanyl)-9-(3-hydroxypropyl)-9H-purin-6-ylamine;** <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.82(m, 2H, CH<sub>2</sub>), 3.60 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 4.12(m, 2H, CH<sub>2</sub>), 4.21 (t, J = 7.0 Hz, 2H, CH<sub>2</sub>), 6.47(s, 1H, Ar-H), 6.86 (d, J = 8.9 Hz,

1H, Ar-H), 7.02 (d,  $J = 8.9$  Hz, 1H, Ar-H); 8.15 (s, 1H, purine-H); electrophile: 1-bromo-3-hydroxypropane.

**Example 12:** The compounds in this example were prepared using diazonium salts and thiols as coupling partners.

### 5 12.1 9-Butyl-8-(2-iodo-5-methoxy-phenylsulfanyl)-9H-purin-6-ylamine

**Step 1:** A suspension of 8-bromo-9-butyl-9H-purin-6-ylamine (0.50 g, 1.85 mmol) and thiourea (1.49 g, 19.6 mmol) in n-butanol (10 ml) was heated to reflux for 14 h. Dilution with  $\text{CH}_2\text{Cl}_2$  (70 ml), washing with water and concentration afforded 6-amino-9-butyl-7,9-dihydro-purine-8-thione as a white powder (0.42 g, 1.87 mmol, 100 %).  $^1\text{H}$

10 NMR ( $\text{DMSO}-d_6$ )  $\delta$  12.35–12.25 (br. s, 1H), 8.13 (s, 1H), 6.92–6.72 (br. s., 2H), 4.09 (t,  $J = 7.6$  Hz, 2H), 1.71 (quint.,  $J = 7.5$  Hz, 2H), 1.29 (sext.,  $J = 7.5$  Hz, 2H), 0.87 (t,  $J = 7.4$  Hz, 3H).

**Step 2:** A solution of the above thione (30.8 mg, 0.138 mmol) and t-BuOK (15.5 mg, 0.138 mmol) in MeOH (0.55 ml) was treated portion-wise with crude 2-iodo-5-methoxy-benzenediazonium tetrafluoroborate (48 mg, 0.138 mmol). The vigorous  $\text{N}_2$  evolution ceased after 2 min. Work-up and preparative TLC ( $\text{MeOH}:\text{CH}_2\text{Cl}_2$  5:95) yielded the title sulfide.

9-Butyl-8-(2-iodo-5-methoxy-benzyl)-9H-purin-6-ylamine;  $R_t = 8.45$  min;  $^1\text{H}$ NMR ( $\text{CDCl}_3-d$ ):  $\delta$  8.38 (s, 1H), 7.73 (d,  $J = 8.7$  Hz, 1H), 6.71 (d,  $J = 2.7$  Hz, 1H), 6.58 (dd,  $J = 8.7$ , 2.7 Hz, 1H), 5.91 (s, 2H), 4.22 (t,  $J = 7.4$  Hz, 2H), 3.68 (s, 3H), 1.75 (quint.,  $J = 7.7$  Hz, 2H), 1.34 (sext.,  $J = 7.5$  Hz, 2H), 0.93 (t,  $J = 7.4$  Hz, 3H).

### **Example 13: Fluorescence-based Competitive Binding Assay for Biotinylated-Geldanamycin to Purified Hsp90**





This assay directly measures the binding of biotinylated-geldanamycin (biotin-GM) to purified Hsp90 and thus tests the ability of compounds to compete for binding to Hsp90.

Purified native Hsp90 protein (**mixture of alpha and beta**) from HeLa cells (Stressgen Biotechnologies Corp., San Diego, CA, USA) was coated onto 96-well plates







by incubating for 1hr at 37°C. Uncoated Hsp90 was removed and the wells washed twice in 1x PBS (phosphate-buffered saline) buffer. Biotin-GM was then added to the wells, and the reaction was further incubated for 1hr 37°C. The wells were washed twice with 1 x PBS, before the addition of 20ug/ml streptavidin-phycoerythrin, and incubated for 1hr at 37°C. The wells were again washed twice with 1x PBS. The fluorescence was then measured in a Gemini spectrofluorometer (Molecular Devices) using an excitation of 485nm and emission of 580nm.







The compounds in Table 4 were synthesized and evaluated for HSP90 binding ability based on the above assay:

Table 4

Example #	IC <sub>50</sub> μM	Example #	IC <sub>50</sub> μM
 <b>1.1</b>	10	 <b>3.4</b>	2.8
 <b>2.4</b>	2.0	 <b>2.2</b>	1.1



Example #	IC <sub>50</sub> μM	Example #	IC <sub>50</sub> μM
 3.2	6	 11.10	1.3
 2.1	2	 4.7	1.1
 4.8	0.9	 4.9	2.3

Example #	IC <sub>50</sub> μM	Example #	IC <sub>50</sub> μM
 <b>9.4</b>	1.5	 <b>4.10</b>	0.9
 <b>9.5</b>	1.8	 <b>9.6</b>	0.9
 <b>11.3</b>	4.0	 <b>9.7</b>	0.8

#### Example 14: HER2 Inhibition Assay

MCF-7 cells are seeded in 24 well plates at a density of approximately 30,000 cells/well and allowed to grow for 16 hours in DMEM supplemented with 10% FBS.



- 5 Drug is then added at a concentration range of 100 μM to .01 μM. Cells are incubated for an additional 24. Drug treated cells and untreated control cells are trypsinized, and







incubated at room temperature for 15 minutes with anti Her-2 neu Ab conjugated with phycoerythrin (Becton Dickinson, San Jose CA; Cat no. 340552) at a concentration of 0.25 ug/ml, or non-specific control IgG1 conjugated with phycoerythrin (Becton Dickinson, San Jose CA; Cat no. 340761). Samples were analyzed using a FACS Calibur flow cytometer (Becton Dickinson) equipped with Argon-ion laser that which emits 15 mW of 488 nm light for excitation of the phycoerythrin fluorochrome. 10,000 events were collected per sample. A fluorescence histogram was generated and the mean fluorescence intensity (mfi) of each sample was determined using Cellquest software. The background was defined as the mfi generated from cells incubated with control IgG, and was subtracted from each sample stained with the HER-2/neu Ab. Percent degradation of Her-2 was calculated as follows:







$$\% \text{Her-2 degradation} = (\text{mfi HER-2 sample}) / (\text{mfi HER-2 untreated cells}) \times 100$$



Table 5 summarizes the Her-2 degradation ability of various compounds of the invention:

**Table 5**

Example #	IC50 μM	Example #	IC50 μM
 <p>1.1</p>	6.0	 <p>3.4</p>	1.5

Example #	IC50 $\mu\text{M}$	Example #	IC50 $\mu\text{M}$
 <b>2.4</b>	1.0	 <b>2.2</b>	0.5
 <b>3.2</b>	1.5	 <b>11.10</b>	0.7
 <b>2.1</b>	0.6	 <b>4.7</b>	1.5

Example #	IC50 $\mu\text{M}$	Example #	IC50 $\mu\text{M}$
 <b>4.8</b>	0.8	 <b>4.9</b>	1.0
 <b>9.4</b>	1.5	 <b>4.10</b>	0.8
 <b>9.5</b>	2.0	 <b>9.6</b>	0.3

Example #	IC <sub>50</sub> μM	Example #	IC <sub>50</sub> μM
 11.3	1.4	 9.7	0.3

Inhibitory Concentration 50 (IC<sub>50</sub>) for this assay is the concentration necessary to degrade 50 % of Her 2 expression (protein).

\* \* \*

5        The foregoing examples are not limiting and merely illustrative of various aspects and embodiments of the present invention. All documents cited herein are indicative of the levels of skill in the art to which the invention pertains and are incorporated by reference herein in their entireties. None, however, is admitted to be prior art.

10       One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The methods and compositions described illustrate preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Certain modifications and other uses will occur to those skilled in the art, and are encompassed within the spirit of the invention, as defined by the scope of the claims.



15       The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and

expressions of excluding any equivalents of the features shown and described, or portions thereof. It is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments, optional features, modifications and variations of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the description and the appended claims.

In addition, where features or aspects of the invention are described in terms of Markush groups or other grouping of alternatives, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group or other group, and exclusions of individual members as appropriate, or by proviso.

Other embodiments are within the following claims.

We claim:





alkoxyalkyl, optionally substituted heterocyclic alkylaminoalkyl, optionally substituted alkylcarbonylaminoalkyl,  $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{NR}^4-(\text{CH}_2)_n$ , optionally substituted alkylcarbonyloxyalkyl  $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{O}-(\text{CH}_2)_n$ , hydroxyalkyl, haloalkyl, perhaloalkyl,  $\text{C}(\text{O})\text{R}^2$ ,  $\text{S}(\text{O})_2\text{R}^2$ ,  $\text{C}(\text{O})\text{NR}^4_2$ , and  $\text{C}(\text{O})\text{OR}^2$ ;

5 wherein Y is selected from the group consisting of H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted alicyclic, optionally substituted araalkyl, optionally substituted aryloxyalkyl, optionally substituted alkoxyalkyl, alkylaminoalkyl, alkylcarbonylaminoalkyl  $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{NR}^4-(\text{CH}_2)_n$ , alkylcarbonyloxyalkyl  $-(\text{CH}_2)_n-$   
 10  $\text{C}(\text{O})-\text{O}-(\text{CH}_2)_n$ , optionally substituted heterocyclic, hydroxyalkyl, haloalkyl, perhaloalkyl,  $\text{C}(\text{O})\text{R}^2$ ,  $\text{S}(\text{O})_2\text{R}^2$ ,  $\text{C}(\text{O})\text{NR}^4_2$ , and  $\text{C}(\text{O})\text{OR}^2$ ;

wherein Z is selected from the group consisting of H, halogen,  $\text{CN}$ ,  $\text{OR}^3$ ,  $\text{SR}^3$ , perhaloalkyl, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted alicyclic, optionally substituted araalkyl, optionally substituted aryloxyalkyl, optionally substituted alkoxyalkyl, optionally substituted heterocyclic,  $\text{C}(\text{O})\text{R}^2$ ,  $-\text{S}(\text{O})_2\text{R}^2$ , guanidine, amidine,  
 15 and  $\text{C}(\text{O})\text{NR}^4_2$ ;

wherein  $\text{R}^1$  is independently selected from H, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heteroalkyl, optionally substituted aryl, optionally substituted heterocyclic,  $\text{C}(\text{O})\text{R}^2$ ,  $-\text{C}(\text{O})\text{OR}^2$ ,  $\text{C}(\text{O})\text{NR}^4_2$ ,  $\text{C}(\text{S})\text{OR}^2$ ,  
 20  $\text{C}(\text{S})\text{NR}^4_2$ ,  $\text{PO}_3\text{R}_4$ ,  $\text{SO}_2\text{R}^2$ ;

wherein  $\text{R}^2$  is independently selected from alkyl, heteroalkyl, cycloalkyl, heterocyclic, heteroaryl, and aryl, all optionally substituted;

wherein  $\text{R}^3$  is independently selected from H, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heteroalkyl, optionally substituted aryl, optionally substituted heterocyclic and  $\text{C}(\text{O})\text{NR}^4_2$ ;  
 25

wherein  $\text{R}^4$  is independently selected from the group consisting of H, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heteroalkyl, optionally substituted aryl, and optionally substituted heterocyclic; and

wherein R<sup>5</sup> is selected from H, OH, and optionally substituted alkyl; and

wherein R<sup>6</sup> is independently selected from H, optionally substituted alkyl, OR<sup>3</sup>, SR<sup>3</sup>, NHR<sup>3</sup>, C(O)N(R<sup>4</sup>)<sub>2</sub>, (O)R<sup>5</sup>, NO<sub>2</sub>, CN, halogen, and S(O)<sub>2</sub>R<sup>2</sup>, and

wherein n is from 1 to 3.

5     2.     The compound, tautomer, or pharmaceutically acceptable salt of claim 1 wherein A is further selected from the group consisting of H, halogen, NR<sup>1</sup><sub>2</sub>, OR<sup>3</sup>, SR<sup>3</sup>, optionally substituted lower alkyl, and C(O)N(R<sup>4</sup>)<sub>2</sub>.

3.     The compound, tautomer, or pharmaceutically acceptable salt of claim 1 wherein Z is further selected from the group consisting of H, halogen, CN, OR<sup>3</sup>, SR<sup>3</sup>, optionally substituted lower perhaloalkyl, optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl, and optionally substituted lower aryl;

10

4.     The compound, tautomer, or pharmaceutically acceptable salt of claim 1 wherein X is further selected from S and S(O) (sulfoxide).

5.     The compound, tautomer, or pharmaceutically acceptable salt of claim 1 wherein Y is further selected from the group consisting of optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl, optionally substituted lower aryl, and optionally substituted lower alicyclic.

15

6.     The compound, tautomer, or pharmaceutically acceptable salt of claim 1 wherein Q is further selected from optionally substituted aryl and optionally substituted heteroaryl.

20     7.     The compound, tautomer, or pharmaceutically acceptable salt of claim 1 wherein X is S;

wherein A is selected from the group consisting of Cl, NH<sub>2</sub>, and CH<sub>3</sub>;


wherein Z is selected from the group consisting of H, F, Cl, Br and CF<sub>3</sub>;

wherein Y is selected from the group consisting of optionally substituted C3-C8 alkyl, optionally substituted C3-C8 alkenyl, optionally substituted C3-C8 alkynyl,

25

optionally substituted C1-C10 lower aryl, and optionally substituted C3-C10 alicyclic; and

wherein Q is selected from the group consisting of optionally substituted aryl optionally substituted heterocyclic, and optionally substituted heteroaryl, e.g.,



5

L = O, S, NR<sup>1</sup>

wherein R<sup>1</sup> is independently selected from H, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heteroalkyl, optionally substituted aryl, optionally substituted heterocyclic, C(O)R<sup>2</sup>, -C(O)OR<sup>2</sup>, C(O)NR<sup>4,2</sup>, C(S)OR<sup>2</sup>,  
 10 C(S)NR<sup>4,2</sup>, SO<sub>2</sub>R<sup>2</sup>, PO<sub>3</sub>R<sup>4</sup>;

wherein R<sup>5</sup> is selected from H, OH, and optionally substituted alkyl; and

wherein R<sup>6</sup> is independently selected from H, optionally substituted alkyl, aryl, OR<sup>3</sup>, SR<sup>3</sup>, NR<sup>3,2</sup>, C(O)R<sup>5</sup>, NO<sub>2</sub>, CN, and halogen; and

wherein R<sup>3</sup> is independently selected from H, optionally substituted alkyl,  
 15 optionally substituted cycloalkyl, optionally substituted heteroalkyl, optionally substituted aryl, optionally substituted heterocyclic and C(O)NR<sup>4,2</sup>.

8. The compound, tautomer, or pharmaceutically acceptable salt of claim 7, wherein A is NH<sub>2</sub>; Z is selected from H, Cl and F; Y is selected from -(CH<sub>2</sub>)<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>3</sub>CCH<sub>3</sub>, -(CH<sub>2</sub>)<sub>4</sub>Br, -(CH<sub>2</sub>)<sub>4</sub>Cl, -(CH<sub>2</sub>)<sub>4</sub>OAc, -(CH<sub>2</sub>)<sub>4</sub>NHEt, -(CH<sub>2</sub>)<sub>4</sub>OH, -(CH<sub>2</sub>)<sub>5</sub>Br, -

(CH<sub>2</sub>)<sub>5</sub>Cl, -(CH<sub>2</sub>)<sub>5</sub>OAc, -(CH<sub>2</sub>)<sub>2</sub>-O<sup>i</sup>Pr, and -(CH<sub>2</sub>)<sub>5</sub>OH; and Q is selected from 2,5-dimethoxyphenyl, 2-iodo-5-methoxyphenyl, 4-iodo-5-methoxyphenyl, 2-iodo-4-fluoro-5-methoxyphenyl, 2-bromo-5-methoxyphenyl, 2-chloro-5-methoxyphenyl, 2-iodo-4-iodo-5-methoxyphenyl, 2-iodo-4-bromo-5-methoxyphenyl, 2-iodo-4-chloro-5-methoxyphenyl, and 2-chloro-3,4,5-trimethoxyphenyl.

9. The compound of claim 1 having name 8-(2,5-dimethoxy-phenylsulfanyl)-2-fluoro-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine, wherein A is NH<sub>2</sub>, Z is F, Y is (CH<sub>2</sub>)<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>, X is S, and Q is 2,5-dimethoxyphenyl; or a pharmaceutically acceptable salt thereof.

10. The compound of claim 1 having name 8-(2,5-dimethoxy-phenylsulfanyl)-2-fluoro-9-(pent-4-ynyl)-9H-purin-6-ylamine, wherein A is NH<sub>2</sub>, Z is F, Y is (CH<sub>2</sub>)<sub>3</sub>CCH, X is S, and Q is 2,5-dimethoxyphenyl; or a pharmaceutically acceptable salt thereof.

11. The compound of claim 1 having name 8-(2,5-dimethoxy-phenylsulfanyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine, wherein A is NH<sub>2</sub>, Z is H, Y is (CH<sub>2</sub>)<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>, X is S, and Q is 2,5-dimethoxyphenyl; or a pharmaceutically acceptable salt thereof.

12. The compound of claim 1 having name 8-(2,5-dimethoxy-phenylsulfanyl)-9-(pent-4-ynyl)-9H-purin-6-ylamine, wherein A is NH<sub>2</sub>, Z is H, Y is (CH<sub>2</sub>)<sub>3</sub>CCH, X is S, and Q is 2,5-dimethoxyphenyl; or a pharmaceutically acceptable salt thereof.

13. The compound of claim 1 having name 8-(2-Iodo-5-methoxy-phenylsulfanyl)-2-fluoro-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine, wherein A is NH<sub>2</sub>, Z is F, Y is (CH<sub>2</sub>)<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>, X is S, and Q is 2-iodo-5-methoxyphenyl; or a pharmaceutically acceptable salt thereof.

14. The compound of claim 1 having name 8-(2-Iodo-5-methoxy-phenylsulfanyl)-2-fluoro-9-(pent-4-ynyl)-9H-purin-6-ylamine, wherein A is NH<sub>2</sub>, Z is F, Y is (CH<sub>2</sub>)<sub>3</sub>CCH, X is S, and Q is 2-iodo-5-methoxyphenyl; or a pharmaceutically acceptable salt thereof.

15. The compound of claim 1 having name 8-(2-Iodo-5-methoxy-phenylsulfanyl)-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine, wherein A is NH<sub>2</sub>, Z is H, Y is

$(\text{CH}_2)_2\text{CH}=\text{C}(\text{CH}_3)_2$ , X is S, and Q is 2-iodo-5-methoxyphenyl; or a pharmaceutically acceptable salt thereof.

16. The compound of claim 1 having name 8-(2-Iodo-5-methoxy-phenylsulfanyl)-9-(pent-4-ynyl)-9H-purin-6-ylamine, wherein A is  $\text{NH}_2$ , Z is H, Y is  $(\text{CH}_2)_3\text{CCH}$ , X is S, and  
5 Q is 2-iodo-5-methoxyphenyl; or a pharmaceutically acceptable salt thereof.

17. The compound of claim 1 having name 8-(2-Iodo-5-methoxy-phenylsulfanyl)-2-chloro-9-(4-methyl-pent-3-enyl)-9H-purin-6-ylamine, wherein A is  $\text{NH}_2$ , Z is Cl, Y is  $(\text{CH}_2)_2\text{CH}=\text{C}(\text{CH}_3)_2$ , X is S, and Q is 2-iodo-5-methoxyphenyl; or a pharmaceutically acceptable salt thereof.

10 18. The compound of claim 1 having name 8-(2-Iodo-5-methoxy-phenylsulfanyl)-2-chloro-9-(pent-4-ynyl)-9H-purin-6-ylamine, wherein A is  $\text{NH}_2$ , Z is Cl, Y is  $(\text{CH}_2)_3\text{CCH}$ , X is S, and Q is 2-iodo-5-methoxyphenyl; or a pharmaceutically acceptable salt thereof.

15 19. A pharmaceutical composition comprising the compound, tautomer, or pharmaceutically acceptable salt of any one of claims 1-18 and one or more pharmaceutical carriers or excipients.

20. A method of inhibiting an HSP90, comprising:

contacting a cell having an HSP90 with a compound, tautomer or pharmaceutically acceptable salt or pharmaceutical composition according to any one of claims 1-18.

21. The method of claim 20 wherein said cell is a mammalian cell.

20 22. The method of claim 20 wherein said mammalian cell is human.

23. The method of claim 20 wherein said contacting occurs in vitro.

24. The method of claim 20 wherein said contacting occurs in vivo.

25. The method of claim 20 wherein said contacting is part of an ex vivo procedure.

26. The method of claim 20 wherein said contacting is accomplished by intravenous  
25 administration to a subject.

27. The method of claim 20 wherein said contacting is accomplished by parenteral administration to a subject.

28. The method of claim 20 wherein said contacting is performed in situ.

29. The method of claim 20 wherein said contacting is part of a therapy directed  
5 against cancer cells.

30. The method of claim 29 wherein said cancer cells are selected from breast cancer cells and melanoma cells.

31. The compound, tautomer or pharmaceutically acceptable salt of any of claims 1-7 having from 0 to 25 substitutions, collectively.

10 32. The compound, tautomer or pharmaceutically acceptable salt of claim 31 having from 0 to 15 substitutions, collectively.

33. The compound, tautomer or pharmaceutically acceptable salt of claim 32 having from 0 to 10 substitutions, collectively.

15 34. The compound, tautomer or pharmaceutically acceptable salt of claim 33 having from 0 to 5 substitutions, collectively.

35. The compound, tautomer or pharmaceutically acceptable salt of claim 31 having from 5 to 15 substitutions, collectively.

36. The compound, tautomer or pharmaceutically acceptable salt of claim 31 having 0 substitutions.

20 37. The compound, tautomer or pharmaceutically acceptable salt of claim 31 having between 0 and 5 substitutions, collectively.

38. A pharmaceutical composition comprising the compound, tautomer, or pharmaceutically acceptable salt of any one of claims 31-37 and one or more pharmaceutical excipients.

25

39. A method of inhibiting an HSP90, comprising:

contacting a cell having an HSP90 with an effective amount of a compound, tautomer, pharmaceutically acceptable salt, or pharmaceutical composition according to any one of claims 31-38.

5 40. The method of claim 39 wherein said cell is a mammalian cell.

41. The method of claim 39 wherein said mammalian cell is human.

42. The method of claim 39 wherein said contacting occurs in vitro.

43. The method of claim 39 wherein said contacting occurs in vivo.

44. The method of claim 39 wherein said contacting is part of an ex vivo procedure.

10 45. The method of claim 39 wherein said contacting is accomplished by intravenous administration to a subject.

46. The method of claim 39 wherein said contacting is accomplished by parenteral administration to a subject.

47. The method of claim 39 wherein said contacting is performed in situ.

15 48. The method of claim 39 wherein said contacting is part of a therapy directed against cancer cells.

49. The method of claim 48 wherein said cancer cells are breast cancer or melanoma cells.

20 50. A complex comprising the compound, tautomer, or pharmaceutically acceptable salt of any of claims 1-18 and 31-37 and at least one other compound.

51. The complex of claim 50 wherein one of said at least one other compound is an HSP90.

52. The complex of claim 51 wherein said HSP90 is human.

53. Use of the compound, tautomer thereof, or pharmaceutically acceptable salt thereof of any of claims 1-18 and 31-37 in a chemotherapy regimen.

54. The use of claim 53 wherein said regimen is part of a combinational therapy that makes use of one or more other agents selected from the group consisting of radioisotopes, antibodies, recombinant products, small molecules, antineoplastic agents, Herceptin, taxol, taxanes and taxane derivatives, gleevac, alkylating agents, anti-metabolites; epidophyllotoxin; an antineoplastic enzyme; a topoisomerase inhibitor; procarbazine; mitoxantrone; platinum coordination complexes; biological response modifiers/growth inhibitors; hormonal/anti-hormonal therapeutic agents and haematopoietic growth factors, anthracycline drugs, vinca drugs, mitomycins, bleomycins, cytotoxic nucleosides, tepothilones, discodermolide, pteridine drugs, diynenes, podophyllotoxins, carminomycin, daunorubicin, aminopterin, methotrexate, methopterin, dichloromethotrexate, mitomycin C, porfiromycin, 5-fluorouracil, 6-mercaptopurine, gemcitabine, cytosine arabinoside, podophyllotoxin, podo-phyllotoxin derivatives, etoposide, etoposide phosphate or teniposide, melphalan, vinblastine, vincristine, leurosine, vindesine, leurosine, paclitaxel, estramustine, carboplatin, cyclophosphamide, bleomycin, gemcitabine, ifosamide, melphalan, hexamethyl melamine, thiotepa, cytarabin, idatrexate, trimetrexate, dacarbazine, L-asparaginase, camptothecin, CPT-11, topotecan, ara-C, bicalutamide, flutamide, leuprolide, pyridobenzoindole derivatives, interferons and interleukins.

55. Use of the compound, tautomer thereof, or pharmaceutical salt thereof of any of claims 1-18 and 31-37 for treating one or more of inflammation, infectious disease, autoimmune disease, and ischemia.

56. The method of claim 20 wherein said contacting is accomplished by oral administration to a subject.

57. The method of claim 20 wherein said contacting is accomplished by topical administration to a subject.

58. The method of claim 48 wherein said cancer cells are melanoma cancer cells.






FIGURE 1





FIGURE 1

(Cont.)

2 / 2